

Product datasheet for **TP503337**

Snf8 (NM_033568) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse SNF8, ESCRT-II complex subunit, homolog (S. cerevisiae) (Snf8), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR203337 protein sequence Red =Cloning site Green =Tags(s)

MHRRGVGAGAIAKKKLAEAKYKERGTVLAEDQLAQM SKQLDMFKTNLEEFASKHKQEIRKNPEFRVQFQD
MCATIGVDPLASGKGFWSEMLGVGDFYYELGVQIIEVCLALKHRNGGLITLLEELHQV LKGRGKFAQDVS
QDDLIRAIKKL KALGTGFGIIPVGGTYLIQSVPAELNMDHTVVLQLAEKNGYVTSEIKTSLKWETERAR
QVLEHLLKEGLAWLDLQAPGEAHYWL PALFTDLYSQEISAEAEKEAFP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	28.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_291046
Locus ID:	27681
UniProt ID:	Q9CZ28
RefSeq Size:	947



[View online »](#)

Cytogenetics: 11 59.24 cM

RefSeq ORF: 777

Synonyms: D11Moh34

Summary: Component of the endosomal sorting complex required for transport II (ESCRT-II), which is required for multivesicular body (MVB) formation and sorting of endosomal cargo proteins into MVBs. The MVB pathway mediates delivery of transmembrane proteins into the lumen of the lysosome for degradation. The ESCRT-II complex is probably involved in the recruitment of the ESCRT-III complex. The ESCRT-II complex may also play a role in transcription regulation by participating in derepression of transcription by RNA polymerase II, possibly via its interaction with ELL. Required for degradation of both endocytosed EGF and EGFR, but not for the EGFR ligand-mediated internalization. Required for the exosomal release of SDCBP, CD63 and syndecan (By similarity).[UniProtKB/Swiss-Prot Function]