

Product datasheet for **TP503503**

Ldlrap1 (NM_145554) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse low density lipoprotein receptor adaptor protein 1 (Ldlrap1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR203503 protein sequence Red =Cloning site Green =Tags(s) MVFSLKYLGMTLVERPKGEELSAAVKRIVATAKASGKKLQKVTCLKVSPRGIIITDLSLTSQLIENVSIYR ISYCTADKMHDKVFAYIAQSQQNESLECHAFLCTKRKVAQAVTLTVAQAFKVAFEFWQVSKEEKEKREKA NQEGGDVPGTRRDSTPSLKTIVATGNLLDLEEVAKAPLSTVSANTNNVDETTPRPQVLGNNNSVWELDDG L DEAFSRLAQSRTPNPQVLDTGLSAQDIHYAQCLSPTDWDKPDSSGIDQDDDVFTF TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	29.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_663529</u>
Locus ID:	100017
UniProt ID:	<u>Q8C142</u>


[View online »](#)

RefSeq Size:	2671
Cytogenetics:	4 D3
RefSeq ORF:	792
Synonyms:	AA691260; Arh; Arh1; ARH2; FHCB1; FHCB2
Summary:	<p>Adapter protein (clathrin-associated sorting protein (CLASP)) required for efficient endocytosis of the LDL receptor (LDLR) in polarized cells such as hepatocytes and lymphocytes, but not in non-polarized cells (fibroblasts). May be required for LDL binding and internalization but not for receptor clustering in coated pits. May facilitate the endocytosis of LDLR and LDLR-LDL complexes from coated pits by stabilizing the interaction between the receptor and the structural components of the pits. May also be involved in the internalization of other LDLR family members. Binds to phosphoinositides, which regulate clathrin bud assembly at the cell surface. Required for trafficking of LRP2 to the endocytic recycling compartment which is necessary for LRP2 proteolysis, releasing a tail fragment which translocates to the nucleus and mediates transcriptional repression (By similarity). [UniProtKB/Swiss-Prot Function]</p>