

Product datasheet for **TP503594**

Prkab1 (NM_031869) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse protein kinase, AMP-activated, beta 1 non-catalytic subunit (Prkab1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR203594 protein sequence Red =Cloning site Green =Tags(s) MGNTSSERAALERQAGHKTPRRDSSGGAKDGRPKILMDSPEDADIFHSEEIKAPEKEEFLAWQHLEAN DKAPAQARPTVFRWTGGGKEVYLSGSFNNWSKLPLTRSQNNFVAILDLPEGEHQYKFFVDGQWTHDPSE P IVTSQLGTVNNIIQVKKTDFEVDALMVDSQKCSDVSELSSPPGPYHQEPYMSKPEERFKAPPILPPHL LQVILNKDTGISCDPALLPEPNHVMLNHLYALSIDKGVMLVLSATHRYKKKYVTLLYKPI TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	30.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_114075</u>
Locus ID:	19079
UniProt ID:	<u>Q9R078</u>


[View online »](#)

RefSeq Size: 2051

Cytogenetics: 5 F

RefSeq ORF: 810

Synonyms: 1300015D22Rik; AU021155; E430008F22

Summary: Non-catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Beta non-catalytic subunit acts as a scaffold on which the AMPK complex assembles, via its C-terminus that bridges alpha (PRKAA1 or PRKAA2) and gamma subunits (PRKAG1, PRKAG2 or PRKAG3) (By similarity).[UniProtKB/Swiss-Prot Function]