

Product datasheet for TP503874

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Bcdin3d (NM_029236) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse BCDIN3 domain containing (Bcdin3d), with C-terminal

MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse Expression Host: HEK293T

Expression cDNA Clone >MR203874 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAADGTLSRGGVGEAVEEEHPGALEPGAAPFGNFPHYSRFHPPEQRLRLLPPELLRQLFPPEGPEKRPIL GLDVGCNSGDLSVALYKHFLSPRDGETCSGASRELRILCCDIDPVLVERAERDCPFPEALTFITLDIMDQ ESRKVPLSSFLSQFGRSVFDMVFCMSVTMWIHLNHGDRGLCEFLAHVSSLCSYLLVEPQPWKCYRAAARR LRKLGLHSFDHFRSLAIRGDMAKQIVRILTQDHGMELACCFGNTSWDRSLLLFRAKHTHETQAIPESSTK

ETRTD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK

Predicted MW: 32 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 083512

 Locus ID:
 75284

 UniProt ID:
 Q91YP1





Bcdin3d (NM_029236) Mouse Recombinant Protein - TP503874

RefSeq Size: 1271
Cytogenetics: 15 F1
RefSeq ORF: 855

Synonyms: 4930556P03Rik; AV138748

Summary: O-methyltransferase that specifically monomethylates 5'-monophosphate of cytoplasmic

histidyl tRNA, acting as a capping enzyme. Less efficiently, also methylates the 5'

monophosphate of pre-miRNAs, acting as a negative regulator of miRNA processing. The 5' monophosphate of pre-miRNAs is recognized by DICER1 and is required for pre-miRNAs processing: methylation at this position reduces the processing of pre-miRNAs by DICER1. Able to mediate methylation of pre-miR-145, as well as other pre-miRNAs. There is some controversy about the methylation of pre-miR-145, since the dimethylation first described as the specific enzymatic activity cannot be reproduced by a more recent work which observes a monomehtylation of pre-miR-145 but two orders weaker than the methylation of cytosolic

histidyl tRNA.[UniProtKB/Swiss-Prot Function]