

Product datasheet for TP504127

Emc2 (NM_025736) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse ER membrane protein complex subunit 2 (Emc2), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR204127 protein sequence Red =Cloning site Green =Tags(s)

MAKVTERYDVTWEEMRDKMRKWREENSRNSEQIMEVGEELINDYASKLGGDDIWIIEQVMIAALDYGRDD
LALFCLQELRRQFPGSHRVKRLTGMRFEAMERYDDAIQLYDRILQEDPTNTAARKRKIAIRKAQGKTVEA
IRELNEYLEQFVGDQEAWHELAEELYINEHDYAKAAFCLEELMMTNPHNHLYCQQYAEVKYTQGGLENLEL
SRKYFAQALKLNNRNMRALFGLYMSASHIASNPKASAKMKKDNIKYASWAANQINRAYQFAGRSKKETKY
SLKAVEDMLETQITQS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	34.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_080012
Locus ID:	66736
UniProt ID:	Q9CRD2 , Q6A0D1



[View online »](#)

RefSeq Size:	1245
Cytogenetics:	15 B3.2
RefSeq ORF:	894
Synonyms:	4921531G14Rik; AV060620; AW209495; Ttc35

Summary: Part of the endoplasmic reticulum membrane protein complex (EMC) that enables the energy-independent insertion into endoplasmic reticulum membranes of newly synthesized membrane proteins. Preferentially accommodates proteins with transmembrane domains that are weakly hydrophobic or contain destabilizing features such as charged and aromatic residues. Involved in the cotranslational insertion of multi-pass membrane proteins in which stop-transfer membrane-anchor sequences become ER membrane spanning helices. It is also required for the post-translational insertion of tail-anchored/TA proteins in endoplasmic reticulum membranes. By mediating the proper cotranslational insertion of N-terminal transmembrane domains in an N-exo topology, with translocated N-terminus in the lumen of the ER, controls the topology of multi-pass membrane proteins like the G protein-coupled receptors. By regulating the insertion of various proteins in membranes, it is indirectly involved in many cellular processes.[UniProtKB/Swiss-Prot Function]