

Product datasheet for TP504246

Cdk4 (NM_009870) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse cyclin-dependent kinase 4 (Cdk4), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR204246 protein sequence Red =Cloning site Green =Tags(s)
	<p>MAATRYEPVAEIGVGAYGTVYKARDPHSGHFVALKSVRVPNGGAAGGGLPVSTVREVALRRLEAFEHPN VRLMDVCATSRTDRDIK/TLVFEHIDQDLRTYLDKAPPPGLPVETIKDLMRQFLSGLDFLHANCIVHRD LKPENILVTSNGTVKLADDFGLARIYSYQMALTPVWVTLWYRAPEVLLQSTYATPVMWSVGCIFAEMFRR KPLFCGNSEADQLGKIFDLIGLPPEDDWPREVSLPRGAFAPRGPRPVQSVPEMEESGAQLLLEMLTFNP HKRISAFRALQHSYLHKEESDAE</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	33.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_034000</u>
Locus ID:	12567
UniProt ID:	<u>P30285</u> , <u>Q545C3</u>



[View online »](#)

RefSeq Size: 1365

Cytogenetics: 10 D3

RefSeq ORF: 912

Synonyms: Crk3

Summary: Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity. Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).[UniProtKB/Swiss-Prot Function]