

## Product datasheet for **TP504285**

### Wdr61 (NM\_001025375) Mouse Recombinant Protein

#### Product data:

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse WD repeat domain 61 (Wdr61), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

**Species:** Mouse

**Expression Host:** HEK293T

**Expression cDNA Clone or AA Sequence:** >MR204285 protein sequence  
Red=Cloning site Green=Tags(s)

MTNQYSILFKQEQAHDDAIWSVAWETNKKENIETVVTGSLDDLKVKWKWRDERLELQWSLEGHQLGVSV  
DISHTLPIAASSSLDAHIRLWDLKQMKSIDAGPVDAWTLAFSPDSQYLATGTHMGKVNIFGVESGKK  
EYSLDTRGKFILSIAYSPDGKYLASGAIDGIINIFDIATGKLLHTLEGHAMP IRSLT FSPDSQLLV T ASD  
DGYIKIYDVQHANLAGT LSGHASWVLNVAFCPDDTHFVSSSSDKSVKVDVGTRTC IHTFFDHQDQVWGV  
KYNGNGSKIVSVGDDQEIHVYDCPI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Tag:** C-MYC/DDK

**Predicted MW:** 33.8 kDa

**Concentration:** >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

**Storage:** Store at -80°C after receiving vials.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** [NP\\_001020546](#)

**Locus ID:** 66317

**UniProt ID:** [Q9ERF3](#), [A0MNP4](#)



[View online »](#)

RefSeq Size:	1331
Cytogenetics:	9 A5.3
RefSeq ORF:	918
Synonyms:	2700038L12Rik; 2810418I05Rik; REC14
Summary:	<p>Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. Required for mono- and trimethylation on histone H3 'Lys-4' (H3K4me3), dimethylation on histone H3 'Lys-79' (H3K4me3). Required for Hox gene transcription. Component of the SKI complex which is thought to be involved in exosome-mediated RNA decay and associates with transcriptionally active genes in a manner dependent on PAF1C (By similarity).[UniProtKB/Swiss-Prot Function]</p>