

Product datasheet for TP504705

Ccnh (NM_023243) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse cyclin H (Ccnh), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR204705 protein sequence Red =Cloning site Green =Tags(s)
	<p>MYHSSSQKRHWTFASEEQLARLRADANRKFCKAVANGKVLNPDPVFLEPHEELTLCKYYEKRLLEFCSV FKPAMPRSVGTACMYFKRFYLNNSVMEYHPRIIMLTCAFLACKVDEFNVSSPQFVGNLRESPLGQERAL EQILEYELLLIQQLNFHLIVHNPYRPFEGFLIDIKTRYPMLENPEILRKTADDFLSRIALTDAYLLYTPS QIALTAILSSASRAGITMESYLSLMLKENRTCLSQLLDIMKSMRNLVKKYEPPRSDEVAVLKQKLERC HSSDLALNAVTKKRKGYEDDDYVSKKPKQEEEEWTDDDLVDSL</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	37.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_075732
Locus ID:	66671
UniProt ID:	Q61458



[View online »](#)

RefSeq Size: 1976

Cytogenetics: 13 C3

RefSeq ORF: 972

Synonyms: 6330408H09Rik; AI661354; AV102684; AW538719

Summary: Regulates CDK7, the catalytic subunit of the CDK-activating kinase (CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation. CAK complexed to the core-TFIIF basal transcription factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminal domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA polymerase II. Its expression and activity are constant throughout the cell cycle.[UniProtKB/Swiss-Prot Function]