

## Product datasheet for TP505005

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Efnb2 (NM 010111) Mouse Recombinant Protein**

**Product data:** 

**Product Type: Recombinant Proteins** 

Description: Purified recombinant protein of Mouse ephrin B2 (Efnb2), with C-terminal MYC/DDK tag,

expressed in HEK293T cells, 20ug

Species: Mouse **Expression Host:** HEK293T

**Expression cDNA Clone** 

>MR205005 protein sequence Red=Cloning site Green=Tags(s) or AA Sequence:

> MAMARSRRDSVWKYCWGLLMVLCRTAISRSIVLEPIYWNSSNSKFLPGQGLVLYPQIGDKLDIICPKVDS KTVGQYEYYKVYMVDKDQADRCTIKKENTPLLNCARPDQDVKFTIKFQEFSPNLWGLEFQKNKDYYIIST SNGSLEGLDNQEGGVCQTRAMKILMKVGQDASSAGSARNHGPTRRPELEAGTNGRSSTTSPFVKPNPGS

TDGNSAGHSGNNLLGSEVALFAGIASGCIIFIVIIITLVVLLLKYRRRHRKHSPQHTTTLSLSTLATPKR

GGNNNGSEPSDVIIPLRTADSVFCPHYEKVSGDYGHPVYIVQEMPPQSPANIYYKV

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

C-MYC/DDK Tag: Predicted MW: 37.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

For testing in cell culture applications, please filter before use. Note that you may experience Note:

some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

NP 034241 RefSeq:

Locus ID: 13642 UniProt ID: P52800





## Efnb2 (NM\_010111) Mouse Recombinant Protein - TP505005

RefSeq Size: 4319

**Cytogenetics:** 8 3.42 cM

RefSeq ORF: 1008

Synonyms: ELF-2; Epl5; Eplg5; Htk-L; LERK-5; Lerk5; NLERK-1

Summary: Cell surface transmembrane ligand for Eph receptors, a family of receptor tyrosine kinases

which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Binds to receptor tyrosine kinase including EPHA4, EPHA3 and EPHB4. Together with EPHB4 plays a central role

in heart morphogenesis and angiogenesis through regulation of cell adhesion and cell migration. EPHB4-mediated forward signaling controls cellular repulsion and segregation from EFNB2-expressing cells. May play a role in constraining the orientation of longitudinally

projecting axons.[UniProtKB/Swiss-Prot Function]