

## **Product datasheet for TP505182**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Slc25a33 (NM\_027460) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse solute carrier family 25, member 33 (Slc25a33), with

C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

>MR205182 protein sequence Red=Cloning site Green=Tags(s)

MATGTQQKENTLLHLFAGGCGGTVGAIFTCPLEVIKTRLQSSRLALRTVYYPQVHLGTISGAGMVRPTSV TPGLLQVLKSILEKEGPKSLFRGLGPNLVGVAPSRAVYFACYSKAKEQFNGVFVPNSNTVHILSAGSAAF VTNTLMNPIWMVKTRMQLERKVRGCKQMNTLQCARRVYQTEGVRGFYRGLTASYAGISETIICFAIYESL KKCLKDAPIVSSTDGAEKSSSGFFGLMAAAAVSKGCASCIAYPHEVIRTRLREEGSKYRSFVQTARLVFR

EEGYLAFYRGLFAQLIRQIPNTAIVLSTYEFIVYLLGERA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK

**Predicted MW:** 35 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 081736

 Locus ID:
 70556

 UniProt ID:
 Q3TZX3





## Slc25a33 (NM\_027460) Mouse Recombinant Protein - TP505182

RefSeq Size: 1877
Cytogenetics: 4 E2
RefSeq ORF: 960

**Synonyms:** 5730438N18Rik; Pnc1

**Summary:** Mitochondrial transporter that imports/exports pyrimidine nucleotides into and from

mitochondria. Transports preferentially uracil, thymine, and cytosine (deoxy)nucleoside diand triphosphates by an antiport mechanism. Also transports guanine but not adenine (deoxy)nucleotides. Is inhibited strongly by pyridoxal 5'-phosphate, 4,7-diphenyl-1,10-phenanthroline, tannic acid, and mercurials (mercury dichloride, mersalyl acid, p-

hydroxymercuribenzoate). Participates in mitochondrial genome maintenance, regulation of mitochondrial membrane potential and mitochondrial respiration (By similarity). Upon INS or IGF1 stimulation regulates cell growth and proliferation by controlling mitochondrial DNA replication and transcription, the ratio of mitochondria-to nuclear-encoded components of

the electron transport chain resulting in control of mitochondrial ROS production (PubMed:17596519). Participates in dendritic cell endocytosis and may associate with mitochondrial oxidative phosphorylation (By similarity).[UniProtKB/Swiss-Prot Function]