

Product datasheet for TP505210

Anxa1 (NM_010730) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse annexin A1 (Anxa1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR205210 protein sequence Red =Cloning site Green =Tags(s)
	<p>MAMVSEFLKQARFLENQEQEYVQAVKSYKGGPGSAVSPYPSFNVSSDVAALHKAIMVKGVDEATIIDILT KRTNAQRQQIKAAYLQENGKPLDEVLRKALTGHLEEVVLAMLKTPAQFDADELRGAMKGLGTDEDTLIEI LTTTRSNEQIREINRVYREELKRDLDITS DTS GDFRKALLALAKGDRCQDLSVNQDLADTDARALYEAG ERRKGTDVNVFTTILTSRSPHLRRVFQNYGKYSQHDMNKALDLELKGDIKCLTTIVKCATSTPAFFAE KLYEAMKGGAGTRHKALIRIMVSRSEIDMNEIKVFYQKKYGISLCQAILDET KGDY EKILVALCGGN</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	38.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_034860
Locus ID:	16952
UniProt ID:	P10107 , Q4FIV4



[View online »](#)

RefSeq Size:	1395
Cytogenetics:	19 13.83 cM
RefSeq ORF:	1041
Synonyms:	Anx-1; Anx-A1; C430014K04Rik; Lpc-1; Lpc1
Summary:	<p>Plays important roles in the innate immune response as effector of glucocorticoid-mediated responses and regulator of the inflammatory process. Has anti-inflammatory activity (PubMed:12475898). Plays a role in glucocorticoid-mediated down-regulation of the early phase of the inflammatory response (PubMed:12475898). Promotes resolution of inflammation and wound healing (PubMed:25664854). Functions at least in part by activating the formyl peptide receptors and downstream signaling cascades. Promotes chemotaxis of granulocytes and monocytes via activation of the formyl peptide receptors (By similarity). Contributes to the adaptive immune response by enhancing signaling cascades that are triggered by T-cell activation, regulates differentiation and proliferation of activated T-cells (PubMed:17948261). Promotes the differentiation of T-cells into Th1 cells and negatively regulates differentiation into Th2 cells (PubMed:17948261). Has no effect on unstimulated T-cells. Promotes rearrangement of the actin cytoskeleton, cell polarization and cell migration. Negatively regulates hormone exocytosis via activation of the formyl peptide receptors and reorganization of the actin cytoskeleton (By similarity). Has high affinity for Ca(2+) and can bind up to eight Ca(2+) ions (By similarity). Displays Ca(2+)-dependent binding to phospholipid membranes (By similarity). Plays a role in the formation of phagocytic cups and phagosomes (PubMed:21245195). Plays a role in phagocytosis by mediating the Ca(2+)-dependent interaction between phagosomes and the actin cytoskeleton (PubMed:21245195).</p> <p>[UniProtKB/Swiss-Prot Function]</p>