

## Product datasheet for **TP506149**

### Map2k1 (NM\_008927) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse mitogen-activated protein kinase kinase 1 (Map2k1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR206149 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	MPKKKPTPIQLNPAPDGSVNGTSSAETNLEALQKKLEELDEQQRKRLEAFLTQKQKVGELKDDDFEK ISELGAGNGGVVFKVSHKPSGLVMARKLIHLEIKPAIRNQIIRELQVLHECNSPYIVGFYGAFYSDGEIS ICMEHMDGGSLDQVLKKAGRIPEQILGKVSIAVIKGLTYLREKHKIMHRDVKPSNILVNSRGEIKLCDFG VSGQLIDSMANSFVGTRSYMSPERLQGTHYSVQSDIWSMGLSLVEMAVGRYPPIPPDAKELELLFGCHVE GDAAETPPRPRTGPRPLSSYGMDSRPPMAIFELLDYIVNEPPPKLP SGVFSLEFQDFVNKCLIKNPAERA DLKQLMVHAFIKRSDAEVDFAGWLCSTIGLNQ PSTPTHAASI
	<b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-MYC/DDK
Predicted MW:	43.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_032953</a>
Locus ID:	26395
UniProt ID:	<a href="#">P31938</a>



[View online »](#)

RefSeq Size:	2387
Cytogenetics:	9 34.55 cM
RefSeq ORF:	1179
Synonyms:	MAPKK1; Mek1; MEKK1; Prkmk1
Summary:	<p>Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2. Both MAP2K1/MEK1 and MAP2K2/MEK2 function specifically in the MAPK/ERK cascade, and catalyze the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in the extracellular signal-regulated kinases MAPK3/ERK1 and MAPK1/ERK2, leading to their activation and further transduction of the signal within the MAPK/ERK cascade. Depending on the cellular context, this pathway mediates diverse biological functions such as cell growth, adhesion, survival and differentiation, predominantly through the regulation of transcription, metabolism and cytoskeletal rearrangements. One target of the MAPK/ERK cascade is peroxisome proliferator-activated receptor gamma (PPARG), a nuclear receptor that promotes differentiation and apoptosis. MAP2K1/MEK1 has been shown to export PPARG from the nucleus. The MAPK/ERK cascade is also involved in the regulation of endosomal dynamics, including lysosome processing and endosome cycling through the perinuclear recycling compartment (PNRC), as well as in the fragmentation of the Golgi apparatus during mitosis.[UniProtKB/Swiss-Prot Function]</p>