

## Product datasheet for TP506264

### Cd3eap (NM\_145822) Mouse Recombinant Protein

#### Product data:

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Purified recombinant protein of Mouse CD3E antigen, epsilon polypeptide associated protein (Cd3eap), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
<b>Species:</b>	Mouse
<b>Expression Host:</b>	HEK293T
<b>Expression cDNA Clone or AA Sequence:</b>	>MR206264 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAGTQACSATRFSCPPHFTEMSPDSEPSRFSLEALTGPDTELWLIQAPADFAPQCLNGRRVPLSGSKTVK  
GKLDGKKHRYRVFTSSPQAREATLLASSEAGGRLTCAPAPSGSLRIMEGPQEYLLSRVPLQLIPTSLPP  
QIPAGLRPRFSAFGGSPVTPGPGSASALRSPTSGKRKSTRKGTDASSDTQEAVNRHGAMEVKALGNLGV  
SVKKRKRYFMQEEMEAKTMEPVVAELPVPSATSSKKRKKSKGTETSQVEHTEPVAQTEPPEGTFLFPTKKR  
KRQKEADGTEEVDGIVADSQPQVIVEAQEETILLSPTKKRRKEKRQNLGMEAEIMGPPGVLMMVTEHSEHGL  
QAEVALVSPKTKKKKKGKRVGETEAALPDDFEPQGALAPSKKKKERGARL

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

<b>Tag:</b>	C-MYC/DDK
<b>Predicted MW:</b>	43.1 kDa
<b>Concentration:</b>	>0.05 µg/µL as determined by microplate BCA method
<b>Purity:</b>	> 80% as determined by SDS-PAGE and Coomassie blue staining
<b>Buffer:</b>	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C after receiving vials.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_665821</a>
<b>Locus ID:</b>	70333
<b>UniProt ID:</b>	<a href="#">Q76KJ5</a>



[View online »](#)

RefSeq Size: 2248

Cytogenetics: 7 A3

RefSeq ORF: 1200

Synonyms: 1190028F09; 2610103M17Rik; ASE-1; Ase1; CAST; PAF49

**Summary:** DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Component of RNA polymerase I which synthesizes ribosomal RNA precursors. Involved in UBTF-activated transcription, presumably at a step following PIC formation.[UniProtKB/Swiss-Prot Function]