

Product datasheet for **TP506830**

Ddx39b (NM_019693) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse DEAD (Asp-Glu-Ala-Asp) box polypeptide 39B (Ddx39b), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR206830 protein sequence Red =Cloning site Green =Tags(s)

MAENDVDNELLDYEDDEVETAAGADGTEAPAKKDVKGSYVSIHSSGFRDFLLKPELLRAIVDCGFEHPSE
VQHECIPQAILGMDVLCQAKSGMGKTAVFVLATLQQLPVTGQVSVLVMCHTRELAFQISKEYERFSKYM
PNVKVAVFFGGLSIKKDEEVLKKNCPHIVGTPGRILALARNKSLNLKHIKHFILDECDKMLEQLDMRRD
VQEIFRMTPEHEKQVMMFSATLSKEIRPVCRKFMQDPMEIFVDDETKLTLLHGLQQYYVKLKDNEKNRKLFD
LLDVLEFNQWIFVKSQRCIALAQLLVEQNFPAAIHRGMPQEERLSRYQQFKDFQRRILVATNLFGRG
MDIERVNIAFNYPEDSDTYLHRVARAGRFGTKGLAITFVSDENDAKILNDVQDRFEVNISELPDEIDI
SSYIEQTR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	49 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_062667</u>
Locus ID:	53817



[View online »](#)

UniProt ID:	Q9Z1N5
RefSeq Size:	1756
Cytogenetics:	17 18.6 cM
RefSeq ORF:	1287
Synonyms:	0610030D10Rik; AI428441; Bat-1; Bat1; Bat1a; D6S81Eh; D17H6S81E; D17H6S81E-1
Summary:	<p>Involved in nuclear export of spliced and unspliced mRNA. Assembling component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NFX1 pathway. May undergo several rounds of ATP hydrolysis during assembly of TREX to drive subsequent loading of components such as ALYREF/THOC and CHTOP onto mRNA. Also associates with pre-mRNA independent of ALYREF/THOC4 and the THO complex. Involved in the nuclear export of intronless mRNA; the ATP-bound form is proposed to recruit export adapter ALYREF/THOC4 to intronless mRNA; its ATPase activity is cooperatively stimulated by RNA and ALYREF/THOC4 and ATP hydrolysis is thought to trigger the dissociation from RNA to allow the association of ALYREF/THOC4 and the NXF1-NXT1 heterodimer. Involved in transcription elongation and genome stability (By similarity). [UniProtKB/Swiss-Prot Function]</p>