

## Product datasheet for **TP507103**

### Eif3e (NM\_008388) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse eukaryotic translation initiation factor 3, subunit E (Eif3e), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR207103 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	<p>MAEYDLTTRIAHFLDRHLVFPILLEFLSVKEIYNEKELLQGKDLLSDTNMVDFAFDVYKNLYSDDIPHAL REKRTTVVAQLKQLQAETEPVKMFEDPETTRQMQSTRDGRMLFDYLADKHGFRQEYLDTRYAKFQYE CGNYSGAAEYLYFFRVLVPATDRNALSSLWGKLASEILMQNWDAAEDLTRKETIDNNSVSSPLQSLQQ RTWLIHWSLFVFFNHPKGRDNIIDFLYQPQYLNAIQTMCPHILRYLTAVITNKDVRKRRQVLKDLVKV IQQESYTYKDPITEFVECLYVNFDFDGAQKKLRECESVLVNDFFLVACLEDFIENARLFIFETFCRIHQ ISINMLADKLNMTPEEAERWIVNLIRNARLDAKIDSKLGHVVMGNNAVSPYQQVIEKTKSLSFERSQMLAM NIEKKLNQNSRSEAPNWTQDSGFY</p> <p><b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b></p>
Tag:	C-MYC/DDK
Predicted MW:	52.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_032414</a></u>
Locus ID:	16341



[View online »](#)

UniProt ID:	<u>P60229</u>
RefSeq Size:	1541
Cytogenetics:	15 16.73 cM
RefSeq ORF:	1335
Synonyms:	48kDa; eIF3-p46; eIF3-p48; Eif3s6; Int6
Summary:	<p>Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA<sup>i</sup> and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression. Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway. May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins.</p> <p>[UniProtKB/Swiss-Prot Function]</p>