

Product datasheet for TP507696

Akt1 (NM_009652) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse thymoma viral proto-oncogene 1 (Akt1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	<p>>MR207696 representing NM_009652</p> <p>Red=Cloning site Green=Tags(s)</p> <p>MNDVAIVKEGWLHKRGEYIKTWRPRYFLLKNDGTFIGYKERPQDVDQRESPLNNFSVAQCQLMKTERPR P NTFIIRCLQWTTVIERTFHVETPEEREWATAIQTVADGLKRQEEETMDFRSGSPSDNSGAEEMEVSLAK PKHRVTMNEFEYKLLGKGTFGKVLVKEKATGRYYAMKILKKEVIVAKDEVAHTLTENRVLQNSRHPFL TALKYSFQTHDRLCFVMEYANGGELFFHLSRERVFSEDRARFYGAEIVSALDYHSEKNVYRDLKLENL MLDKDGHKIDTDFGLCKEGIKDGATMKTFCGTPEYLAPEVLEDNDYGRAVDWWGLGVVMYEMMCGRLP FY NQDHEKLFELILMEEIRFPRTLGPPEAKSLLSGLLKKDPTQRLGGGSEDAKEIMQHRFFANIVWQDVYEKK LSPFPKPQVTSETDTRYFDEEFTAQMITITPPDQDDSMECVDSERRPHFPQFSYSASGTA</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	56.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.


[View online »](#)

RefSeq: [NP_033782](#)

Locus ID: 11651

UniProt ID: [P31750](#)

RefSeq Size: 2707


Cytogenetics: 12 61.2 cM

RefSeq ORF: 1440

Synonyms: Ak; Akt; LTR-akt; PK; PKB; PKB/A; PKB/Akt; PKBalpha; Rac

Summary: This gene encodes the founding member of the Akt serine-threonine protein kinase gene family that also includes Akt2 and Akt3. This kinase is a major downstream effector of the phosphatidylinositol 3-kinase (PI3K) pathway that mediates the effects of various growth factors such as platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin and insulin-like growth factor I (IGF-I). It is activated through recruitment to cellular membranes by PI3K lipid products and by phosphorylation by 3-phosphoinositide dependent kinase-1. It then further phosphorylates different downstream proteins in response to various extracellular signals and thus plays a pivotal role in mediating a variety of cellular processes, such as glucose metabolism, glycogen biosynthesis, protein synthesis and turn over, inflammatory response, cell survival (anti-apoptosis) and development. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009]

Product images:



Purified recombinant protein Akt1 was analyzed by SDS-PAGE gel and Coomossie Blue Staining.