

Product datasheet for **TP507727**

Hdac1 (NM_008228) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse histone deacetylase 1 (Hdac1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR207727 representing NM_008228 Red =Cloning site Green =Tags(s)
	<p>MAQTQGTQRKVCYYYDGDVGNYYYGQGHMPKPHRIMTHNLLLNLYGLYRKMEIYRPHKANAEEMTKY HSD DYIKFLRSIRPDNMSEYSKQMQRFNVEDCPVFDGLFEFCQLSTGGSVASAVKLNKQQTDAVNWAGGL H HAKKSEASGFCYVNDIVLAILELLKYHQRVLYIDIDIHHDGVEEAFYTTDRVMTVSFHKYGEYFPGTGD LRDIGAGKGKYYAVNYPLRDGIDDESIEAIFKPVMSKVMEMFQPSAVVLQCGSDSLSGDRLGCFNLTIKG HAKCVEFVKSFNLPMLMLGGGGYTIRNVARCWYETAVALDTEIPNELPYNDYFEYFGPDFKLHISPSNM TNQNTNEYLEKIKQRLFENLRMLPHAPGVQMQAIPEDAIPESGDEDEEDPDKRISICSSDKRIACEEEF SDSDEEGEGGRKNSSNFKKAKRVKTEDEKEKDPEEKKEVTEEEKTKEEKPEAKGVKEEVKLA</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	55.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq:	<u>NP_032254</u>
Locus ID:	433759
UniProt ID:	<u>O09106</u>
RefSeq Size:	1971
Cytogenetics:	4 63.26 cM
RefSeq ORF:	1446
Synonyms:	HD1; Hdac1-ps; MommeD5; RPD3
Summary:	<p>Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Deacetylates SP proteins, SP1 and SP3, and regulates their function. Component of the BRG1-RB1-HDAC1 complex, which negatively regulates the CREST-mediated transcription in resting neurons. Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation. Deacetylates TSHZ3 and regulates its transcriptional repressor activity. Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B. Deacetylates NR1D2 and abrogates the effect of KAT5-mediated relieving of NR1D2 transcription repression activity. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. Involved in CIART-mediated transcriptional repression of the circadian transcriptional activator: CLOCK-ARNTL/BMAL1 heterodimer. Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex or CRY1 through histone deacetylation.[UniProtKB/Swiss-Prot Function]</p>