

Product datasheet for **TP508113**

Prpf19 (NM_134129) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse pre-mRNA processing factor 19 (Prpf19), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR208113 representing NM_134129 Red =Cloning site Green =Tags(s)
	<p>MSLICISISNEVPEHPCVSPVSNHVVYERRLIEKYIAENGTDPINNQPLSEEQOLIDIKVAHPIRPKPPSATS IPAILKALQDEWDAVMLHSFTLRQQLQTTTRQELSHALYQHDAACRVIARLTKEVTAAREALATLKPQAGL IVPQAVPSSQPSVVGAGEPMDLGELVGMTPEIIQKLQDKATVLTTERKKRGKTVPEELVKPEELSKYRQV ASHVGLHSASIPGILALDLCPSDTNKILTGGADKNVVFDFKSTEQILATLKGHTKKVTSVVFHPSQELVF SASPDATIRIWSVPNTSCVQVWRAHESAVTGLSLHATGDYLLSSDDQYWAFSDIQTGRVLTKVTDETSG CSLTCAQFHPDGLIFGTGTMDSQIKIWDLKERTNVANFPGHSGPITSIAFSENGYYLATAADSSVKLWD LRKLKNFKTLQLDNNFEVKSLIFDQSGTYLALGGTDVQIYICKQWTEILHFTHEHSLTTGVAFGHHAKFI ASTGMDRSLKFYSL</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	55.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_598890</u>



[View online »](#)

Locus ID:	28000
UniProt ID:	Q99KP6
RefSeq Size:	6161
Cytogenetics:	19 7.33 cM
RefSeq ORF:	1512
Synonyms:	AA617263; AL024362; D19Wsu55e; NMP200; Prp19; PSO4; Snev
Summary:	<p>Isoform 1: Ubiquitin-protein ligase which is a core component of several complexes mainly involved in pre-mRNA splicing and DNA repair. Required for pre-mRNA splicing as component of the spliceosome. Core component of the PRP19C/Prp19 complex/NTC/Nineteen complex which is part of the spliceosome and participates in its assembly, its remodeling and is required for its activity. During assembly of the spliceosome, mediates 'Lys-63'-linked polyubiquitination of the U4 spliceosomal protein PRPF3. Ubiquitination of PRPF3 allows its recognition by the U5 component PRPF8 and stabilizes the U4/U5/U6 tri-snRNP spliceosomal complex. Recruited to RNA polymerase II C-terminal domain (CTD) and the pre-mRNA, it may also couple the transcriptional and spliceosomal machineries. The XAB2 complex, which contains PRPF19, is also involved in pre-mRNA splicing, transcription and transcription-coupled repair. Beside its role in pre-mRNA splicing PRPF19, as part of the PRP19-CDC5L complex, plays a role in the DNA damage response/DDR. It is recruited to the sites of DNA damage by the RPA complex where PRPF19 directly ubiquitinates RPA1 and RPA2. 'Lys-63'-linked polyubiquitination of the RPA complex allows the recruitment of the ATR-ATRIP complex and the activation of ATR, a master regulator of the DNA damage response. May also play a role in DNA double-strand break (DSB) repair by recruiting the repair factor SETMAR to altered DNA. As part of the PSO4 complex may also be involved in the DNA interstrand cross-links/ICLs repair process. In addition, may also mediate 'Lys-48'-linked polyubiquitination of substrates and play a role in proteasomal degradation (PubMed:17349974). May play a role in the biogenesis of lipid droplets (PubMed:17118936). May play a role in neural differentiation possibly through its function as part of the spliceosome (By similarity).[UniProtKB/Swiss-Prot Function]</p>