

Product datasheet for **TP508439**

Slc3a2 (NM_008577) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse solute carrier family 3 (activators of dibasic and neutral amino acid transport), member 2 (Slc3a2), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR208439 protein sequence Red =Cloning site Green =Tags(s)

MSQDTEVDMKDVLENELEPEKQPMNAADGAAAGEKNGLVKIKVAEDETEAGVKFTGLSKEELLKVAGSPG
WVRTRWALLLLFWLGWLGMLAGAVIIVRAPRCREL PVQRWWHKGALYRIGDLQAFVGRDAGGIAGLKSH
LEYLSTLKVKGLVLGPIHKNQKDEINETDLKQINPTLGSQEDFKDLLQSAKKKSIHIILDLPNYQGQNA
WFLPAQADIVATKMKEALSSWLQDGVDFQFRDVGKLMNAPLYLAEWQNITKNLSEDRLLIAGTESSDLQ
QIVNILESTSDLLLSSYLSNSTFTGERTESLVTRFLNATGSQWCSWSVSQAGLLADFI PDHLLRLYQLL
LFTLPGTVPVFSYGDELGLQGALPGQPAKAPLMPWNESSIFHIPPVSLNMTVKQONEDPGSLLTQFRRLS
DLRGKERSLLHGDFHALSSSPDLFSYIRHWDQNERYLVVLNFRDSSGRSARLGASNLPAGISLPASAKLLL
STDSARQSREEDTSLKLENLSLNPYEGLLLQFPFVA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	58.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_032603](#)

Locus ID: 17254

UniProt ID: [P10852](#)

RefSeq Size: 2572

Cytogenetics: 19 5.44 cM

RefSeq ORF: 1581

Synonyms: 4F2; 4F2HC; AI314110; Cd98; Ly-10; Ly-m10; Ly10; Mdu1; Mgp-2hc; NACAE

Summary: Component of several heterodimeric amino acid transporter complexes. The precise substrate specificity depends on the other subunit in the heterodimer (PubMed:9915839). The heterodimer with SLC3A2 functions as sodium-independent, high-affinity transporter that mediates uptake of large neutral amino acids such as phenylalanine, tyrosine, L-DOPA, leucine, histidine, methionine and tryptophan (PubMed:9915839). The complexes with SLC7A6 and SLC7A7 mediate uptake of dibasic amino acids. The complexes function as amino acid exchangers (By similarity). Required for targeting of SLC7A5 and SLC7A8 to the plasma membrane and for channel activity (PubMed:9915839). Plays a role in nitric oxide synthesis in human umbilical vein endothelial cells (HUVECs) via transport of L-arginine (By similarity). The heterodimer with SLC7A5/LAT1 may play a role in the transport of L-DOPA across the blood-brain barrier (Probable). May mediate blood-to-retina L-leucine transport across the inner blood-retinal barrier (By similarity). The heterodimer with SLC7A5/LAT1 can mediate the transport of thyroid hormones triiodothyronine (T3) and thyroxine (T4) across the cell membrane. When associated with SLC7A5 or SLC7A8, involved in the cellular activity of small molecular weight nitrosothiols, via the stereoselective transport of L-nitrosocysteine (L-CNSO) across the transmembrane. The heterodimer with SLC7A5 is involved in the uptake of toxic methylmercury (MeHg) when administered as the L-cysteine or D,L-homocysteine complexes. Together with ICAM1, regulates the transport activity SLC7A8 in polarized intestinal cells, by generating and delivering intracellular signals. When associated with LAPT4B, the heterodimer formed by SLC3A2 and SLC7A5 is recruited to lysosomes to promote leucine uptake into these organelles, and thereby mediates mTORC1 activation (By similarity). [UniProtKB/Swiss-Prot Function]