

## Product datasheet for **TP508579**

### Ehd3 (NM\_020578) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse EH-domain containing 3 (Ehd3), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR208579 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	<p>MFSWLGNDRRKKDPEVFQTVSDGLKKLYTKLLPLEEYRFHEFHSPALEDADFDNKPMVLLVGQYSTG KTFIRYLLEQDFPGMRIGPEPTTDSFIAMQGDVEGIIPGNALVDPKPKPFRKLNFAFNLRVCAQ LPNAVLESISVIDTPGILSGEKQRISRGYDFAAVLEWFAERVDRIILLFDAHKLDISDEFSEVIKALKNH EDKMRVVLNKADQIETQQLMRVYGALMWSLGKIVNTPEVIRVYIGSFWSHPLLIPDNRKLFEEEQDLFR DIQSLPRNAALRKLNDLIKRRARLAKVHAYIISLKKEMPSVFGKDTKKKELVNNLAEIYGRIEREHQISP GDFPNLKRMQDQLQAQDFSKFQPLKSKLLEVDDMLAHDIAQLMVLVRQEETQRPVQMVKGGAFEGTLQG PFGHGYGEGAGEGIDDAEWWWARDKPMYDEIFYTLSPVDGKITGANAKKEMVRSKLPNSVLGKIWLADI DKDGMLDDEEFALANHLIKVKLEGHELPSLPAHLLPPSKRKVSE</p> <p><b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b></p>
Tag:	C-MYC/DDK
Predicted MW:	60.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_065603</a></u>



[View online »](#)

Locus ID: 57440

UniProt ID: [Q9QXY6](#)

RefSeq Size: 3658

Cytogenetics: 17 45.2 cM

RefSeq ORF: 1608

Synonyms: Ehd2

**Summary:** ATP- and membrane-binding protein that controls membrane reorganization/tubulation upon ATP hydrolysis. In vitro causes tubulation of endocytic membranes (By similarity). Binding to phosphatidic acid induces its membrane tubulation activity (PubMed:26896729). Plays a role in endocytic transport. Involved in early endosome to recycling endosome compartment (ERC), retrograde early endosome to Golgi, and endosome to plasma membrane (rapid recycling) protein transport. Involved in the regulation of Golgi maintenance and morphology (By similarity). Involved in the recycling of internalized D1 dopamine receptor (By similarity). Plays a role in cardiac protein trafficking probably implicating ANK2. Involved in the ventricular membrane targeting of SLC8A1 and CACNA1C and probably the atrial membrane localization of CACNA1GG and CACNA1H implicated in the regulation of atrial myocyte excitability and cardiac conduction (PubMed:20489164, PubMed:24759929, PubMed:25825486). In conjunction with EHD4 may be involved in endocytic trafficking of KDR/VEGFR2 implicated in control of glomerular function (PubMed:21408024). Involved in the rapid recycling of integrin beta-3 implicated in cell adhesion maintenance (By similarity). Involved in the unidirectional retrograde dendritic transport of endocytosed BACE1 and in efficient sorting of BACE1 to axons implicating a function in neuronal APP processing. Plays a role in the formation of the ciliary vesicle, an early step in cilium biogenesis; possibly sharing redundant functions with Ehd1 (PubMed:25686250). [UniProtKB/Swiss-Prot Function]