

Product datasheet for TP509054

Msn (NM_010833) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins
Description: Purified recombinant protein of Mouse moesin (Msn), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species: Mouse
Expression Host: HEK293T
Expression cDNA Clone or AA Sequence: >MR209054 representing NM_010833
Red=Cloning site **Green**=Tags(s)

MPKTISVRVTMDAELEFAIQPNTTGKQLFDQVVKITGLREVWFFGLQYQDTKAFSTWLKLNKKVTAQDV
RKESPLLFKFRAKFYPEDVSEELIQDITQRLFFLQVKEGILNDDIYCPPETAVLLASYAVQSKYGDFNKE
VHKSGYLAGDKLLPQRVLEQHKLKNDQWEERIQVWHEEHRGMLREDAVLEYLKIAQDLEMYGVNYFSIKN
KKGSELWLGVDALGLNIYEQNDRLTPKIGFPWSEIRNISFNDKKFKVFKIPDKKAPDFVYAPRLRINKRI
LALCMGNHELYMRRRKPDITIEVQQMKAQAREEKHQKQMERALLENEKKKRELAEEKEKIEREKEELMEK
LKQIEEQTKKAQQELEEQTRRALELEQERKRAQSEAEKLAKERQEAEEAKEALLQASRDQKKTQEQLASE
MAELTARISQLEMARKKKESEAVEWQQAQMVQEDLEKTRAELKTAMSTPHVAEPAENEHDEQDENGAEA
SAELRADAMAKDRSEERTTEAEKNERVQKHLKALTSELANARDESKKTANDMIHAENMRLGRDKYKTLR
QIRQGNTKQRIDEFESM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK
Predicted MW: 68.2 kDa
Concentration: >0.05 µg/µL as determined by microplate BCA method
Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage: Store at -80°C after receiving vials.
Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_034963](#)
Locus ID: 17698
UniProt ID: [P26041](#)
RefSeq Size: 3840
Cytogenetics: X C3
RefSeq ORF: 1731
Synonyms: C78546

Summary: Ezrin-radixin-moesin (ERM) family protein that connects the actin cytoskeleton to the plasma membrane and thereby regulates the structure and function of specific domains of the cell cortex. Tethers actin filaments by oscillating between a resting and an activated state providing transient interactions between moesin and the actin cytoskeleton (By similarity). Once phosphorylated on its C-terminal threonine, moesin is activated leading to interaction with F-actin and cytoskeletal rearrangement (By similarity). These rearrangements regulate many cellular processes, including cell shape determination, membrane transport, and signal transduction (By similarity). The role of moesin is particularly important in immunity acting on both T and B-cells homeostasis and self-tolerance, regulating lymphocyte egress from lymphoid organs (PubMed:22875842). Modulates phagolysosomal biogenesis in macrophages (PubMed:28978692). Participates also in immunologic synapse formation (By similarity). [UniProtKB/Swiss-Prot Function]