

Product datasheet for TP509493

Cdc16 (NM_027276) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse CDC16 cell division cycle 16 (Cdc16), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR209493 representing NM_027276 Red=Cloning site Green=Tags(s)

MNLEPLRKRVRQYLDQQYQSALFWADKVASLSHEEPQDVYWLAQCLYLTAQYHRAAAHALRSRKLDKLYE
ACRYLAARCHYAAKEHQALDILDMEEPINRRLFEKYLKDDNGSRDPSSDWEMSQSSIKSSICLLRGKIY
DALDNRTLATYSYKEALKLDVYCFAFDLLTSHHMLTAQEEKELDSLPLNKLCAEEQELLRFVFENKLLK
KYNKPSETVIPESVDGLQENLDVWVSLAERHYNCDFKMCYKLTSTVMEKDPFHANCLPVIHIGTLVELNK
ANELFYLSHKLVLDLYPSNPVSWFAVGCYMLVGHKNEHARRYLSKATTLEKTYGPAWIAYGHSFAVESEH
DQAMAAYFTAAQLMKGCHLPMLYIGLEYGLTNNKLAERFFGQALSAPEDPFIHEVGVAFQNGEWKT
AEKWFLDALEKIKAGNEVTVDKWEPLLNGLGHVCRKLKKYAEALDYHRQALVLIPQNASTYSAIGYIHS
LMGNFENAVDYFHTALGLRRDDTFSVTMLGHCIEMYIGDSEAYIGADIKDKLKCDFDVHTMKTLKNIIS
PPWDFRDFEVEKQNTTEAGLAPLQNSTKAPESRPNLEETFEIEMNESDMMLETSMDSHST

SGPTRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	71.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq:	NP_081552
Locus ID:	69957
UniProt ID:	Q8R349 , Q3TI84
RefSeq Size:	2253
Cytogenetics:	8 A1.1
RefSeq ORF:	1860
Synonyms:	2700071J12Rik; 2810431D22Rik; APC6
Summary:	Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (By similarity). [UniProtKB/Swiss-Prot Function]