

## Product datasheet for TP509591

### Syk (NM\_011518) Mouse Recombinant Protein

#### Product data:

|                                       |  |
|---------------------------------------|--|
| Product Type:                         | Recombinant Proteins   |
| Description:                          | Purified recombinant protein of Mouse spleen tyrosine kinase (Syk), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug  |
| Species:                              | Mouse  |
| Expression Host:                      | HEK293T  |
| Expression cDNA Clone or AA Sequence: | >MR209591 protein sequence<br><span style="color: red;">Red</span> =Cloning site <span style="color: green;">Green</span> =Tags(s) |

MAGSAVDSANHLTYLFGNITREEAEDYLVQGGMTDGLYLLRQSRNYLGGFALSVAHNRKAHHYTIERELN  
 GTYAISGGRAHASPADLCHYHSQEPDGLICLLKKPFNRPPGVQPKTGFEDLKENLIREYVKQTWNLQGG  
 ALEQAIISQKPQLEKLIATTAHEKMPWFHGNISRDESEQTVLIGSKTNGKFLIRARDNSGSYALCLLHEG  
 KVLHYRIDRDKTGKLSIPEGKKFDTLWQLVEHYSYKPDGLLRVLTVPCKIGAQMHPGSPNAHPVTWSP  
 GGIISRIKSYSPKPGHKKPAPPQGSRPSTVSFNPYEPTGGPWGPDRGLQREALPMDTEVYESPYADPE  
 EIRPKEVYLDRLTLEDNELGSGNFGTVKKGYQMKKVVKTVAVKILKNEANDPALKDELLAEANVMQQ  
 LDNPYIVRMIGICEAESWMLVMEMAELGPLNKYLQQNRHIKDKNIIELVHQVSMGMKYLEESNFVHRDLA  
 ARNVLLVTQHYAKISDFGLSKALRADENYYKAQTHGKWPVKWYAPECINYYKFSSKSDVWSFGVLMWEAF  
 SYGQKPYRGMKGSEVTAMLEKGERMGCPAGCPREMYDLMLNLCWTYDVENRPGFTAVELRLRNYYYDV  
 N

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

|                |  |
|----------------|--|
| Tag:           | C-MYC/DDK  |
| Predicted MW:  | 71.3 kDa   |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method   |
| Purity:        | > 80% as determined by SDS-PAGE and Coomassie blue staining  |
| Buffer:        | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol   |
| Note:          | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. |
| Storage:       | Store at -80°C after receiving vials.  |
| Stability:     | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.        |


[View online »](#)

|               |                  |
|---------------|------------------|
| RefSeq:       | <u>NP_035648</u> |
| Locus ID:     | 20963            |
| UniProt ID:   | <u>P48025</u>    |
| RefSeq Size:  | 5148             |
| Cytogenetics: | 13 27.41 cM      |
| RefSeq ORF:   | 1887             |
| Synonyms:     | Sykb             |

**Summary:**

Non-receptor tyrosine kinase which mediates signal transduction downstream of a variety of transmembrane receptors including classical immunoreceptors like the B-cell receptor (BCR). Regulates several biological processes including innate and adaptive immunity, cell adhesion, osteoclast maturation, platelet activation and vascular development. Assembles into signaling complexes with activated receptors at the plasma membrane via interaction between its SH2 domains and the receptor tyrosine-phosphorylated ITAM domains. The association with the receptor can also be indirect and mediated by adapter proteins containing ITAM or partial hemITAM domains. The phosphorylation of the ITAM domains is generally mediated by SRC subfamily kinases upon engagement of the receptor. More rarely signal transduction via SYK could be ITAM-independent. Direct downstream effectors phosphorylated by SYK include VAV1, PLCG1, PI-3-kinase, LCP2 and BLNK. Initially identified as essential in B-cell receptor (BCR) signaling, it is necessary for the maturation of B-cells most probably at the pro-B to pre-B transition. Activated upon BCR engagement, it phosphorylates and activates BLNK an adapter linking the activated BCR to downstream signaling adapters and effectors. It also phosphorylates and activates PLCG1 and the PKC signaling pathway. It also phosphorylates BTK and regulates its activity in B-cell antigen receptor (BCR)-coupled signaling. In addition to its function downstream of BCR plays also a role in T-cell receptor signaling. Plays also a crucial role in the innate immune response to fungal, bacterial and viral pathogens. It is for instance activated by the membrane lectin CLEC7A. Upon stimulation by fungal proteins, CLEC7A together with SYK activates immune cells inducing the production of ROS. Also activates the inflammasome and NF-kappa-B-mediated transcription of chemokines and cytokines in presence of pathogens. Regulates neutrophil degranulation and phagocytosis through activation of the MAPK signaling cascade. Required for the stimulation of neutrophil phagocytosis by IL15 (By similarity). Also mediates the activation of dendritic cells by cell necrosis stimuli. Also involved in mast cells activation. Involved in interleukin-3/IL3-mediated signaling pathway in basophils (PubMed:19098920). Also functions downstream of receptors mediating cell adhesion. Relays for instance, integrin-mediated neutrophils and macrophages activation and P-selectin receptor/SELPG-mediated recruitment of leukocytes to inflammatory loci. Plays also a role in non-immune processes. It is for instance involved in vascular development where it may regulate blood and lymphatic vascular separation. It is also required for osteoclast development and function. Functions in the activation of platelets by collagen, mediating PLCG2 phosphorylation and activation. May be coupled to the collagen receptor by the ITAM domain-containing FCER1G. Also activated by the membrane lectin CLEC1B that is required for activation of platelets by PDPN/podoplanin. Involved in platelet adhesion being activated by ITGB3 engaged by fibrinogen. Together with CEACAM20, enhances production of the cytokine CXCL8/IL-8 via the NFkB pathway and may thus have a role in the intestinal immune response (PubMed:26195794).[UniProtKB/Swiss-Prot Function]