

Product datasheet for TP510090

App (NM_007471) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse amyloid beta (A4) precursor protein (App), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR210090 protein sequence Red=Cloning site Green=Tags(s)

MLPSLALLLLAAWTVRALEVPTDGNAGLLAEPQIAMFCGKLNMHMNVQNGKWESDPSGKTCTCIGTKEGIL
QYCQEVYPQLQITNVVEANQPVTIQNWCKRGRKQCKTHTHIVIPYRCLVGEFVSDALLVPDKCKFLHQR
MDVCETHLHWHTVAKETCSEKSTNLHDYGMLLPCGIDKFRGVEFVCCPLAEESDVSADAEEDSDVWWW
GGADTDYADGGEDKVVEVAEEEEVADVEEEEADDEDVEDGDEVEEEAEEPYEATERTTSTATTTTTTT
ESVEEVRVPTTAASTPDAVDKYLETPGDENEHAHFQKAKERLEAKHRERMSQVMREWEAERQAKNLPK
ADKKAVIQHFQEKVESLEQEAANERQQLVETHMARVEAMLNDRRRLALENYITALQAVPPRPHHVFNMLK
KYVRAEQKDRQHTLKHFEHVRMVDPKKAAQIRSQVMTHLRVIYERMNQSLSLYNVPAVAEEIQDEVDEL
LQKEQNYSDDLANMISEPRISYGNLALMPSLTETKTTVELLPVNGEFLDDLPWHPFGVDSVPANTEN
EVEPVDARPAADRGLTTRPGSGLTNIKTEEISEVKMDAEFGHDSGFVVRHQKLVFFAEDVGSNKGAIIGL
MVGGVVIATVIVITLVMLKKKQYTSIHHGVVEVDAAVTPEERHLSKMQQNGYENPTYKFFEQMQN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-MYC/DDK
Predicted MW:	78.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq:	NP_031497
Locus ID:	11820
UniProt ID:	Q6GR78
RefSeq Size:	3152
Cytogenetics:	16 46.92 cM
RefSeq ORF:	2088
Synonyms:	Abeta; Abpp; Adap; Ag; betaApp; Cvap; E030013M08Rik
Summary:	<p>Functions as a cell surface receptor and performs physiological functions on the surface of neurons relevant to neurite growth, neuronal adhesion and axonogenesis. Interaction between APP molecules on neighboring cells promotes synaptogenesis. Involved in cell mobility and transcription regulation through protein-protein interactions. Can promote transcription activation through binding to APBB1-KAT5 and inhibit Notch signaling through interaction with Numb. Couples to apoptosis-inducing pathways such as those mediated by G(O) and JIP. Inhibits G(o) alpha ATPase activity (By similarity). Acts as a kinesin I membrane receptor, mediating the axonal transport of beta-secretase and presenilin 1. May be involved in copper homeostasis/oxidative stress through copper ion reduction. Can regulate neurite outgrowth through binding to components of the extracellular matrix such as heparin and collagen I and IV (By similarity). The splice isoforms that contain the BPTI domain possess protease inhibitor activity. Induces a AGER-dependent pathway that involves activation of p38 MAPK, resulting in internalization of amyloid-beta peptide and leading to mitochondrial dysfunction in cultured cortical neurons (By similarity). Provides Cu(2+) ions for GPC1 which are required for release of nitric oxide (NO) and subsequent degradation of the heparan sulfate chains on GPC1. [UniProtKB/Swiss-Prot Function]</p>

Product images:



Purified recombinant protein App was analyzed by SDS-PAGE gel and Coomassie Blue Staining.