

Product datasheet for **TP510392**

Ddx1 (NM_134040) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse DEAD (Asp-Glu-Ala-Asp) box polypeptide 1 (Ddx1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA	>MR210392 protein sequence
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MAAFSEMGMPEIAQAVEEMDWLLPTDIQAESIPLILGGGDVLMMAETGSGKTGAFSIPVIQIVYETLKD
QQEGKKGKTTIKTGASVLNKWQMNPYDRGSAFAIGSDGLCCQSREVKWHGCRGTRGLLKGKHYYEVSCH
DQGLCRVWSTMQASLDLGTDFGFGGGTGKKSHNKQFDNYGEEFTMHDTIGCYLDIDKGHVKFSKNGK
DLGLAFEIPAHIKNQALFPACVLKNAELKFNFGEEEFKFPKDG FVALSKAPDNYIVKSQHTGNAQVSQT
KFLPNAPKALIVEPSRELAEQTLNNVKQFKKYIDNPKLRELLIIGGVAARDQLSVLDNGVDIVGTPGRL
DDLVTGKLNLSQVRFLVLEADGLLSQGYSDFINRMHNQIPQITCDGKRLQVIVCSATLHSFDVKKLSE
KIMHFPTWVDLKGEDSVPDTHHHVVPVNPKTDKLWERLGNHIRTDDVHAKDNTRPGANSPMWSEAIK
ILKGEYAVRAIKEHKMDQAIIFCRTKIDCDNLEQYFMQQGGPDKKGGHGFSCVCLHGD RPKPHERKQNLER
FKKGDVRFLECTDVAARGIDIHGVPYVINVTLPDEKQNYVHRIGRVGRAERMGLAISLVATEKEKVVYHV
CSNRGKGCYNTRLKEDGGCTIWNEMQLLSEIEEHLNCTISQVEPDIKVPVDFDGDGKVTYGGQKRAAGGGN
YKGHVDVLAPTVQELAALKEAQTSLHLGYLPNQLFRFTF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	82.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_598801
Locus ID:	104721
UniProt ID:	Q91VR5
RefSeq Size:	2488
Cytogenetics:	12 A1.1
RefSeq ORF:	2223
Synonyms:	AA409185; DBP-RB
Summary:	Acts as an ATP-dependent RNA helicase, able to unwind both RNA-RNA and RNA-DNA duplexes. Possesses 5' single-stranded RNA overhang nuclease activity. Possesses ATPase activity on various RNA, but not DNA polynucleotides. May play a role in RNA clearance at DNA double-strand breaks (DSBs), thereby facilitating the template-guided repair of transcriptionally active regions of the genome. Together with RELA, acts as a coactivator to enhance NF-kappa-B-mediated transcriptional activation (By similarity). Acts as a positive transcriptional regulator of cyclin CCND2 expression (PubMed:19398953). Binds to the cyclin CCND2 promoter region (PubMed:19398953). Associates with chromatin at the NF-kappa-B promoter region via association with RELA. Binds to poly(A) RNA. May be involved in 3'-end cleavage and polyadenylation of pre-mRNAs. Component of the tRNA-splicing ligase complex required to facilitate the enzymatic turnover of catalytic subunit RTCB: together with archease (ZBTB8OS), acts by facilitating the guanylation of RTCB, a key intermediate step in tRNA ligation (By similarity). Component of a multi-helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the induction of proinflammatory cytokines via the adapter molecule TICAM1 (PubMed:21703541). Specifically binds (via helicase ATP-binding domain) on both short and long poly(I:C) dsRNA (PubMed:21703541).[UniProtKB/Swiss-Prot Function]