

Product datasheet for **TP510582**

Tnfaip3 (NM_009397) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse tumor necrosis factor, alpha-induced protein 3 (Tnfaip3), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >MR210582 protein sequence
 Red=Cloning site Green=Tags(s)

MAEQLLPQALYLSNMRKAVKIRERTPEDIFKPTNGIYHFKTMHRYTLEMFRTCQFCPQFREIIHKALID
 RSVQASLESQKKLNWCREVRLVALKTNGDGNCLMHAACQYMWGVQDSDLVLRKALCSTLKETDTRNF
 KF
 RWQLESLSQEFVETGLCYDTRNWNDEWDNLVKMASADTPAARSGLQYNSLEEIHIFVLSNILRRPIVI
 SDKMLRSLESQSNFAPLKVGGIYLPWHWPAQECYRPIVLGYDSQHFVPLVTLKDSGPELRAVPLVNRDR
 GRFEDLKVHFLTDPENEMKEKLLKEYLIVMEIPVQGWHDGTTHLINAAKLDEANLPKEINLVDDYFELVQ
 HEYKKWQENSQARRAAHAQNPLEPSTPQLSLMDIKCETPNCPFFMSVNTQPLFHECSERRQKNQSKL
 PK
 LNSKLGPEGLPGVGLGSSNWSPEETAGGPHSAPPTAPSLFLFSETTAMKCRSPGCPFTLNVQHNGFCERC
 HARQINASHTADPGKCQACLQDVTRTFNGICSTCFKRTTAEPSSSLTSSIPASCHQRSKSDPSQLIQSLT
 PHSCHRTGNVSPSGCLSQAARTPGDRAGTSKCRKAGCMYFGTPENKGFTLCFIEYRENKQSVTASEKAG
 SPAPRFQNNVPCLGRECGTLGSTMFEGYCQKCFIEAQNRHFHEARRTEEQLRSSQHRDMPRTTQVASRL
 K
 CARASCKNILACRSEELCMQCQHLSQLRVGSVAHRGEPTPEEPPKQRCRAPACDHFGNAKCNNGYCNECYQ
 F
 KQMYG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK

Predicted MW: 88.1 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol



[View online »](#)

Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_033423</u>
Locus ID:	21929
UniProt ID:	<u>Q60769</u>
RefSeq Size:	4437
Cytogenetics:	10 8.08 cM
RefSeq ORF:	2325
Synonyms:	A20; Tnfip3
Summary:	<p>Ubiquitin-editing enzyme that contains both ubiquitin ligase and deubiquitinase activities. Involved in immune and inflammatory responses signaled by cytokines, such as TNF-alpha and IL-1 beta, or pathogens via Toll-like receptors (TLRs) through terminating NF-kappa-B activity. Essential component of a ubiquitin-editing protein complex, comprising also RNF11, ITCH and TAX1BP1, that ensures the transient nature of inflammatory signaling pathways. In cooperation with TAX1BP1 promotes disassembly of E2-E3 ubiquitin protein ligase complexes in IL-1R and TNFR-1 pathways; affected are at least E3 ligases TRAF6, TRAF2 and BIRC2, and E2 ubiquitin-conjugating enzymes UBE2N and UBE2D3. In cooperation with TAX1BP1 promotes ubiquitination of UBE2N and proteasomal degradation of UBE2N and UBE2D3. Upon TNF stimulation, deubiquitinates 'Lys-63'-polyubiquitin chains on RIPK1 and catalyzes the formation of 'Lys-48'-polyubiquitin chains. This leads to RIPK1 proteasomal degradation and consequently termination of the TNF- or LPS-mediated activation of NF-kappa-B. Deubiquitinates TRAF6 probably acting on 'Lys-63'-linked polyubiquitin. Upon T-cell receptor (TCR)-mediated T-cell activation, deubiquitinates 'Lys-63'-polyubiquitin chains on MALT1 thereby mediating disassociation of the CBM (CARD11:BCL10:MALT1) and IKK complexes and preventing sustained IKK activation. Deubiquitinates NEMO/IKBKG; the function is facilitated by TNIP1 and leads to inhibition of NF-kappa-B activation. Upon stimulation by bacterial peptidoglycans, probably deubiquitinates RIPK2. Can also inhibit I-kappa-B-kinase (IKK) through a non-catalytic mechanism which involves polyubiquitin; polyubiquitin promotes association with IKBKG and prevents IKK MAP3K7-mediated phosphorylation. Targets TRAF2 for lysosomal degradation. In vitro able to deubiquitinate 'Lys-11'-, 'Lys-48'- and 'Lys-63' polyubiquitin chains. Inhibitor of programmed cell death. Has a role in the function of the lymphoid system. Required for LPS-induced production of proinflammatory cytokines and IFN beta in LPS-tolerized macrophages.[UniProtKB/Swiss-Prot Function]</p>