

Product datasheet for TP510747

Eif3b (NM_133916) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse eukaryotic translation initiation factor 3, subunit B (Eif3b), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR210747 protein sequence Red=Cloning site Green=Tags(s)
	<p>MQDAENVAVPEAAEERAEPARQQPASESPPTDEAAGSGGSEVGQTEDAEEDAEGPEPEVRAKPAAQSEE ETATSPAASPTPQSAERSPSQEPSAPGKAEAVGEQARGHPSAGAE EGGSDGSAAEAEPRALENGEADEP SFSDPEDFVDDVSEELLGDLVKDRPQADGIDSIVVDNVPQVGPDRLEKLNVIHKIFSKFGKIINDY YPEEDGKTKGYIFLEYASPAHAVDAVKNADGYKLDKQHTFRVNLFTDFDKYMTISDEWDIPEKQPFKDLG NLRWLEEAECRDQYSVIFESGDRTSIFWNDVKDPVSIERARWTETYVRWSPKGTYLATFFHQRIALWG GDKFKQIQRFHQGVQLIDFSPCERYLVTFSPMLMDTQDDPQAIWIDILTGHKKRGFHCESAHWPFIKW SHDGKFFARMTLDTLSIYETPSMGLLDKKSLSKISGIKDFSWSPGGNIAFWVPEDKDIPARVTLMQLPTR QEIRVRNLFNVVDCCLHWQKNGDYLCVKVDRTPKGTQGVVTFEIFRMREKQVPVDVEMKETIIAFAWE PNGSKFAVLHGEAPRISVSFYHVKSNGKIELIKMFDKQANTIFWSPQGQFVLAGLRSMNGALAFVDTS DCTVMNIAEHYMASDVEWDPTGRYVWTSVSWWSHKVDNAYWLWTFQGRLLQKNNKDRFCQLLWRPRPPTL LSQDQIKQIKKDLKKYSKIFEQKDRLSQSKASKELVERRRTMMEDFRQYRMAQELYMKQKNERLELRGG VDTDELDSNVDDWEETIEFFVTEEVIPLGSQE</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	91.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_598677
Locus ID:	27979
UniProt ID:	Q8JZQ9
RefSeq Size:	2966
Cytogenetics:	5 78.97 cM
RefSeq ORF:	2412
Synonyms:	AL033316; AL033334; AL033369; AW208965; D5Wsu45e; EIF3-ETA; EIF3-P110; EIF3-P116; Eif3s9; PRT1
Summary:	<p>RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAⁱ and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression. [UniProtKB/Swiss-Prot Function]</p>