

Product datasheet for TP511094

Nfkb2 (NM_001177370) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse nuclear factor of kappa light polypeptide gene enhancer in B cells 2, p49/p100 (Nfkb2), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >MR211094 representing NM_001177370
Red=Cloning site **Green**=Tags(s)

MDNCYDPGLDGIPEYDDFEFSPSIVEPKDPAPETADGPYLVIVEQPKQRGFRFRYGCESPHGGGLPGASS
EKGRKTYPTVKICNYEGPAKIEVDLVTHSDPPRAHAHSLVGKQCSELGVCAVSVGPKDMTAQFNNLGV LH
VTKKNMMEIMIQLQRQLRSKPQGLTEAERRELEQEAKELKKVMDLSIVLRFSAFLRASDGSFSLPLK
PVISQPIHDSKSPGASNLKISRMDKTAGSVRGGDEVYLLCDKVQKDDIEVRFYEDDENGWQAFGDFSPTD
VHKQYAIVFRTPPYHKMKIERPVTVFLQLKRKRGGDVSDSKQFTYYPLVEDKEEVQRKRRKALPTFSQPF
GGGSHMGGGSGGSAGGYGGAGGGGSLGFFSSSLAYNPYQSGAAPMGCYPGGGGGAQMAGSRRDTDAGEGA
EERTPPEAPQGEPQALDQLQRAREYNARLFLGLAQRSARALLDYGVTADARALLAGQRHLLMAQDENGDT
PLHLAIHGGQTGVIEQIAHVIYHAQYLGVINLTNHLHQTPLHLAVITGQTRVVSFLLQVGADPTLLDRHG
DSALHLALRAGAAPELLQALLRSGAHAVPQILHMPDFEGLYPVHLAVHARSPECLDLLVDCGAEVEAPE
RQGGRTALHLATEMEELGLVTHLVTKLHANVNARTFAGNTPLHLAAGLSPTLRLLLKAGADIHAENEE
PLCPLPSPSTSGSDSDSEGPERDTQRNFRGHTPLDLTCSTKVKTLNNAQNTTEPLAPPSPAGPGLSL
GDAALQNLEQLLDGPEAQGSWAELAERLGLRSLVDTYRKTPSPSGSLLRSYKLAGGDLVGLLEALSMDMGL
HEGVRLKGPETRDKLPSTEVKEDSAYGSQSVEQAEKLCPPPEPPGGLCHGHPQPQVH

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK

Predicted MW: 97.4 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining


Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.



[View online »](#)

Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_001170841</u>
Locus ID:	18034
UniProt ID:	<u>Q9WTK5</u>
RefSeq Size:	2973
Cytogenetics:	19 38.8 cM
RefSeq ORF:	2697
Synonyms:	lyt; NF-kappaB2; p49; p49/p100; p50B; p52
Summary:	<p>NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. In a non-canonical activation pathway, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. The NF-kappa-B heterodimeric RelB-p52 complex is a transcriptional activator. The NF-kappa-B p52-p52 homodimer is a transcriptional repressor. NFKB2 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p100 and generation of p52 by a cotranslational processing. The proteasome-mediated process ensures the production of both p52 and p100 and preserves their independent function. p52 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. p52 and p100 are respectively the minor and major form; the processing of p100 being relatively poor. Isoform p49 is a subunit of the NF-kappa-B protein complex, which stimulates the HIV enhancer in synergy with p65 (By similarity). In concert with RELB, regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-ARNTL/BMAL1 heterodimer.[UniProtKB/Swiss-Prot Function]</p>

Product images:

Purified recombinant protein Nfkb2 was analyzed by SDS-PAGE gel and Coomassie Blue Staining.