

Product datasheet for **TP514387**

Ccdc62 (NM_001134767) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins
Description: Purified recombinant protein of Mouse coiled-coil domain containing 62 (Ccdc62), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species: Mouse
Expression Host: HEK293T
Expression cDNA Clone or AA Sequence: >MR214387 representing NM_001134767
Red=Cloning site **Green**=Tags(s)

MRSSEGAPSWAVALPPPLRPCAYGVSEVTRCWHQLSLGAGESSMNPSTLYRRQNIGSEVETSTIEKQRK
ELQLLIGELKDRDKELNDMVAHVHQRQLLSWEEDRQKVLTLLEERCSKLEGELHKRDTIISLMKKVKTLES
NQAECQTALQKTQQQLQEMAQKATHSTLLSEDLARNENLSSTLVDLSAQVQGLQAREQALTTMIKLDK
DII EAVNHISDCSGKFKLLEHALRDAKMAETCVVREKQDYKQKALRIEVNKLKEDLNEKTENNEQRE
EIIRLKQEKSLHDELIFTVEREKRKDELLEIAKSKQDRTNSELQNLRQIYVKQQSDLQFLNFNISSQE
LIQIHGLKMEEPKALECSKDMCLSDLDNNYPKIDIKRERNQKSLVKDQTFEVMLAQHNGSDKSSCDACRE
KKLQVNTALGEKSVALSSLFTKDLLDKQKSWSLGGKIQTENKVTLCVHAKSPKCDGVLPTTEEKQL
SETSVLSLDEKQWHDINVYGLSSCSKQPDRLDGDGHDRTGTSEVSCCTPNVVICIGDNDLSESKCCHPSN
IIIEAPGHMTDTEWMNIFKPSRAQRIVRHKTMCTCSRSVSAMKYNSSASELIGMQPSQCVGSLKSAEREE
ESAALPDRRTSANEKDDFSPTSKLQRLLAESRQMVTDLELSTLLPISCENLNRSKLEVSEEPDEKTTLV
H

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK
Predicted MW: 79.3 kDa
Concentration: >0.05 µg/µL as determined by microplate BCA method
Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage: Store at -80°C after receiving vials.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001128239
Locus ID:	208908
UniProt ID:	E9PVD1
RefSeq Size:	3920
Cytogenetics:	5 F
RefSeq ORF:	2103
Synonyms:	AI661708; G1-485-3; Gm150; repro29
Summary:	Nuclear receptor coactivator that can enhance preferentially estrogen receptors ESR1 and ESR2 transactivation. Modulates also progesterone/PGR, glucocorticoid/NR3C1 and androgen/AR receptors transactivation, although at lower level; little effect on vitamin D receptor/VDR (By similarity).[UniProtKB/Swiss-Prot Function]