

Product datasheet for TP516559

Anapc7 (NM_019805) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse anaphase promoting complex subunit 7 (Anapc7), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR216559 protein sequence Red =Cloning site Green =Tags(s)
	<p>MSVIDHVRDMAAAGLHSNVRLSSLLLTMSNNNPELFSPSQKYQLLVYHADSLFHDKEYRNAVSKYAMAL QQKKALSKTSKVRPSTGNSASTPQSQCPLPSEIEVKYKMAECYTMLKLDKDAIAVLGDGIPSRQRTPKINMM LANLYKKAGQERPSVTSYKEVLRQCPLALDAILGLLSLVKGAEVASMTMNVIQTPNLDWLSVWIKAYA FVHTGDNSRAINTICSLEKKSLLRDNDVLLGSLADLYFRAGDSKNSVLKFEQAQMLDPYLIRGMDVYGYL LAREGRLEDVENLGCRLFNISDQHAEPWVWVGCHSFYSKRYSRALYLGAKAIQLNSNSVQALLLKGAALR NMGRVQEAIHFREAIRLAPCRLDCYEGLIECYLASNSIREAMVMANNVYKTLGANAQTLTLLATVCLED PVTQEAKATLLDKALAQRPDYVKAVVKAELLSREQKYEDGIALLRNALANQSDCVLHRLGDFLAVNE YQEAMDQYSIALSLDPNDQKSLEGMQKMEEESPTDATQEEDVDDMEGSGEEDLEGSDSEAAQWADQEQ WFGMQ</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	63 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_062779](#)

Locus ID: 56317

UniProt ID: [Q9WVM3](#)

RefSeq Size: 2751

Cytogenetics: 5 F

RefSeq ORF: 1698

Synonyms: APC7; AW545589

Summary: Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (By similarity).[UniProtKB/Swiss-Prot Function]