

## Product datasheet for TP519078

### Armc8 (NM\_028768) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse armadillo repeat containing 8 (Armc8), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR219078 representing NM_028768 Red=Cloning site Green=Tags(s)

MACLLETPIRMSVLSEVTASSRHYVDRLFDPDPQKVLQGVDMKNAVIGNNKQKANLIVLGAVPRLLYLL  
QQETSSTELKTECAVVLGSLAMGTENNPKSLLDCHIIPALLQGLLSPDLKFIEACLRCLRTIFTSPVTP  
ELLYTDATVIPHLMALLSRSTYQEQYICQIFSHCKGPDHQTILFNHGAVQNIHLLTSPSYKVRMQALK  
CFSVLAFENPQVSMTLVNVLDGELLPQIFVKMLQRDKPIEMQLTSAKCLTYMCRAGAIRTDSCIVLKT  
LPCLVRMCSKERLLEERVEGAETLAYLIEPDVELQRIASITDHLIAMLADYFKYPSSVSAITDIKRLDHD  
LKHAEHLRQAQAFKLYASLGANDEDIRKKIETETMMDRIVTGLSESSVKVRLAAVRLHSLSRVQQLRT  
SFQDHAVWKPMLKVLQNPDEILVASSMLCNLLLEFSPSKEPILES GAVELLCGLTQSEN PALRVNGI  
WALMNMAFQAEQKIKADILRSLSTEQLFRLLSDSDMNVLMKTLGLLRNLSTRPHIDKIMSTHGKQIMQAV  
TLILEGEHSIEVKEQTLCILANIADGTTAKELIMTNDILQKIKYYMGHSHVKLQLAAMFCISNLIWNEE  
EGSQRQDKLRDMGIVDILHKLQSADS NLCDKAKTALQQYLA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	75.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq:	<a href="#">NP_083044</a>
Locus ID:	74125
UniProt ID:	<a href="#">Q9DBR3</a> , <a href="#">G3X920</a>
RefSeq Size:	4628
Cytogenetics:	9 E3.3
RefSeq ORF:	2019
Synonyms:	1200015K23Rik; Gid5; HSPC056
Summary:	Component of the CTLH E3 ubiquitin-protein ligase complex that selectively accepts ubiquitin from UBE2H and mediates ubiquitination and subsequent proteasomal degradation of the transcription factor HBP1.[UniProtKB/Swiss-Prot Function]