

Product datasheet for TP521165

Cenpx (NM_016665) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse centromere protein X (Cenpx), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	A DNA sequence from Mouse cDNA ORF Clone, MR221165, encoding Mouse full-length Cenpx.
Tag:	C-MYC/DDK
Predicted MW:	9.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_057874
Locus ID:	20892
UniProt ID:	Q8C4X1
RefSeq Size:	655
Cytogenetics:	11 E2
RefSeq ORF:	234
Synonyms:	Stra13


[View online »](#)

Summary:

DNA-binding component of the Fanconi anemia (FA) core complex. Required for the normal activation of the FA pathway, leading to monoubiquitination of the FANCI-FANCD2 complex in response to DNA damage, cellular resistance to DNA cross-linking drugs, and prevention of chromosomal breakage. In complex with CENPS (MHF heterodimer), crucial cofactor for FANCM in both binding and ATP-dependent remodeling of DNA. Stabilizes FANCM. In complex with CENPS and FANCM (but not other FANC proteins), rapidly recruited to blocked forks and promotes gene conversion at blocked replication forks. In complex with CENPS, CENPT and CENPW (CENP-T-W-S-X heterotetramer), involved in the formation of a functional kinetochore outer plate, which is essential for kinetochore-microtubule attachment and faithful mitotic progression. As a component of MHF and CENP-T-W-S-X complexes, binds DNA and bends it to form a nucleosome-like structure. DNA-binding function is fulfilled in the presence of CENPS, with the following preference for DNA substates: Holliday junction > double-stranded > splay arm > single-stranded. Does not bind DNA on its own. [UniProtKB/Swiss-Prot Function]