

## **Product datasheet for TP522043**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Psmd13 (NM\_011875) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse proteasome (prosome, macropain) 26S subunit, non-

ATPase, 13 (Psmd13), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

**Species:** Mouse

**Expression Host:** HEK293T

**Expression cDNA Clone** >MR222043 representing NM\_011875

or AA Sequence: Red=Cloning site Green=Tags(s)

MKDVPAFLQQSQSSGPGQAAVWHRLEELYTKKLWHQLTLEVLDFVQDPCFAQGDGLIKLYENFISEFEHR VNPLSLVEIILHVVRQMTDPNVALTFLEKTREKVKSSDEAVILCKTAIGALKLNIGDLQATKETIEDVEE MLNNLPGVTSVHSRFYDLSSKYYQTIGNHASYYKDALRFLGCVDIKDLPVSEQQERAFTLGLAGLLGEGV FNFGELLMHPVLESLRDTDRQWLIDTLYAFNSGAVDRFQTLKCAWGQQPDLAANEAQLLRKIQLLCLMEM TFTRPANHRQLTFEEIAKSAKITVNKVELLVMKALSVGLVRGSIDEVDKRVHMTWVQPRVLDLQQIKGMK

DRLELWCTDVKSMEMLVEHQAQDILT

**SGPTRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-MYC/DDK

Predicted MW: 42.8 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 036005

**Locus ID:** 23997

UniProt ID: Q9WV|2, B2RT97





## Psmd13 (NM\_011875) Mouse Recombinant Protein - TP522043

RefSeq Size: 1563

Cytogenetics: 7 F5
RefSeq ORF: 1128
Synonyms: S11

**Summary:** Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent

degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression,

apoptosis, or DNA damage repair.[UniProtKB/Swiss-Prot Function]