

## Product datasheet for TP522290

### Prkcg (NM\_011102) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse protein kinase C, gamma (Prkcg), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR222290 representing NM_011102 Red=Cloning site Green=Tags(s)

MAGLPGGGDSEGGPRPLFCRKGALRQKVHEVKSHKFTARFFKQPTFCSHCTDFIWGIGKQGLQCQVCS  
FVHRRCHEFVTFECPGAGKGPQTDDPRNKHKFRLHSYSSPTFCDHCGSLLYGLVHQGMKSCCEMNVHR  
RCVRSVPSLCGVDHTERRGRLQLEIRAPTSDEIHITVGEARNLIPMDPNGLSDPYVKLKLIPDPRNLTKQ  
KTCTVKATLNPVWNETFVFNLPKGDVERRLSVEVWDWDRTRSNDMFMGAMSGVSELLKAPVDGWYKLLNQ  
EEGEYYNVPVADADNCSLLQKFEACNYPLELYERVMGPSSSPIPSPSPTDSKRCFFGASPGRLHISD  
FSFLMVLGKGSFGKVMLAERRGSDELYAIKILKKDVIQDDDDVDCTLVEKRVLALGGRGPGGRPHFLTQL  
HSTFQTPDRLYFVMEYVTGGDLMIYHIQQLGKFKEPHAAFYAAEIAIGLFFLHNQGIYRDLKLDNVMMLDA  
EGHIKITDFGMCKENVPFGSTTRTFCGTPDYIAPEIAYQPYGKSVDWWSFGVLLYEMLAGQPPFDGEDE  
EELFQAIMEQTVTPKSLSREAVAICKGFLTKHPGKRLGSGPDGEPTIRAHGFFRWIDWERLERLEIAPP  
FRPRPCGRSGENFDKFFTRAAPALTPDRLVLASIDQADFQGFYVNPFDVHPDARSPTSPVPVPM

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	78.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP\\_035232](#)

Locus ID: 18752

UniProt ID: [P63318](#), [Q3UN66](#)

RefSeq Size: 3131

Cytogenetics: 7 1.93 cM

RefSeq ORF: 2091

Synonyms: Pkcc; PKCgamma; Prkcc

**Summary:** Calcium-activated, phospholipid- and diacylglycerol (DAG)-dependent serine/threonine-protein kinase that plays diverse roles in neuronal cells and eye tissues, such as regulation of the neuronal receptors GRIA4/GLUR4 and GRIN1/NMDAR1, modulation of receptors and neuronal functions related to sensitivity to opiates, pain and alcohol, mediation of synaptic function and cell survival after ischemia, and inhibition of gap junction activity after oxidative stress. Binds and phosphorylates GRIA4/GLUR4 glutamate receptor and regulates its function by increasing plasma membrane-associated GRIA4 expression. In primary cerebellar neurons treated with the agonist 3,5-dihydroxyphenylglycine, functions downstream of the metabotropic glutamate receptor GRM5/MGLUR5 and phosphorylates GRIN1/NMDAR1 receptor which plays a key role in synaptic plasticity, synaptogenesis, excitotoxicity, memory acquisition and learning. May be involved in the regulation of hippocampal long-term potentiation (LTP), but may be not necessary for the process of synaptic plasticity. May be involved in desensitization of mu-type opioid receptor-mediated G-protein activation in the spinal cord, and may be critical for the development and/or maintenance of morphine-induced reinforcing effects in the limbic forebrain. May modulate the functionality of mu-type-opioid receptors by participating in a signaling pathway which leads to the phosphorylation and degradation of opioid receptors. May also contribute to chronic morphine-induced changes in nociceptive processing. Plays a role in neuropathic pain mechanisms and contributes to the maintenance of the allodynia pain produced by peripheral inflammation. Plays an important role in initial sensitivity and tolerance to ethanol, by mediating the behavioral effects of ethanol as well as the effects of this drug on the GABA(A) receptors. During and after cerebral ischemia modulate neurotransmission and cell survival in synaptic membranes, and is involved in insulin-induced inhibition of necrosis, an important mechanism for minimizing ischemic injury. Required for the elimination of multiple climbing fibers during innervation of Purkinje cells in developing cerebellum. Is activated in lens epithelial cells upon hydrogen peroxide treatment, and phosphorylates connexin-43 (GJA1/CX43), resulting in disassembly of GJA1 gap junction plaques and inhibition of gap junction activity which could provide a protective effect against oxidative stress. Phosphorylates p53/TP53 and promotes p53/TP53-dependent apoptosis in response to DNA damage. Involved in the phase resetting of the cerebral cortex circadian clock during temporally restricted feeding. Stabilizes the core clock component ARNTL/BMAL1 by interfering with its ubiquitination, thus suppressing its degradation, resulting in phase resetting of the cerebral cortex clock (PubMed:23185022).[UniProtKB/Swiss-Prot Function]