

Product datasheet for TP522510

Pdcd6 (NM_011051) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse programmed cell death 6 (Pdcd6), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR222510 representing NM_011051 Red =Cloning site Green =Tags(s) MAAYSYPGPGGGPGPAAGAALPDQSFLWNVFQRVDKDRSGVISDNELQQALSNGTWTFPNPVTVRSII S MFDRENKAGVNFSEFTGVWKYITDWQNVFRTYDRDNSGMIDKNELKQALSGFGYRLSDQFHDILIRKFD R QGRGQIAFDDFIQGCIVLQRLTDIFRRYDTDQDGWIQVSYEQYLSMVFSIV TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	21.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_035181
Locus ID:	18570
UniProt ID:	P12815


[View online »](#)

RefSeq Size:	1524
Cytogenetics:	13 C1
RefSeq ORF:	573
Synonyms:	alg-2; Alg2; AV299538; MA-3; PS2
Summary:	<p>Calcium sensor that plays a key role in processes such as endoplasmic reticulum (ER)-Golgi vesicular transport, endosomal biogenesis or membrane repair (PubMed:10744743, PubMed:11525164, PubMed:27541325). Acts as an adapter that bridges unrelated proteins or stabilizes weak protein-protein complexes in response to calcium: calcium-binding triggers exposure of apolar surface, promoting interaction with different sets of proteins thanks to 3 different hydrophobic pockets, leading to translocation to membranes (PubMed:10744743, PubMed:11525164, PubMed:27541325). Involved in ER-Golgi transport by promoting the association between PDCD6IP and TSG101, thereby bridging together the ESCRT-III and ESCRT-I complexes (PubMed:10744743, PubMed:11525164, PubMed:27541325). Together with PEF1, acts as calcium-dependent adapter for the BCR(KLHL12) complex, a complex involved in ER-Golgi transport by regulating the size of COPII coats (By similarity). In response to cytosolic calcium increase, the heterodimer formed with PEF1 interacts with, and bridges together the BCR(KLHL12) complex and SEC31 (SEC31A or SEC31B), promoting monoubiquitination of SEC31 and subsequent collagen export, which is required for neural crest specification (By similarity). Involved in the regulation of the distribution and function of MCOLN1 in the endosomal pathway (By similarity). Promotes localization and polymerization of TFG at endoplasmic reticulum exit site (By similarity). Required for T-cell receptor-, Fas-, and glucocorticoid-induced apoptosis (PubMed:8560270). May mediate Ca(2+)-regulated signals along the death pathway: interaction with DAPK1 can accelerate apoptotic cell death by increasing caspase-3 activity (By similarity). Its role in apoptosis may however be indirect, as suggested by knockout experiments (PubMed:12024023). May inhibit KDR/VEGFR2-dependent angiogenesis; the function involves inhibition of VEGF-induced phosphorylation of the Akt signaling pathway (By similarity).[UniProtKB/Swiss-Prot Function]</p>