

## Product datasheet for TP522732

### Cadm1 (NM\_207675) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse cell adhesion molecule 1 (Cadm1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR222732 representing NM_207675 Red=Cloning site Green=Tags(s)

MASAVLPSGSQCAAAA AVAAAAA PGLRLRLLLLLLSAAALPTGDGQNLFTKDVTVIEGEVATISCQVN  
KSDDSVIQLLNPNRQTIYFRDFRPLKDSRFQLLNFSSSELKVSLTNVISIDEGRYFCQLYTDPPQESYTT  
ITVLVPPRNLMIDIQKDTAVEGEEIEVNCTAMASKPATTIRWFKGNKELKGKSEVEEWSDMYTVTSQLML  
KVHKEDDGVPVICQVEHPAVTGNLQTQRYLEVQYKPVVHIQMTYPLQGLTREGDAFELTCEAIGKPQPM  
VTWVRVDDEMPQHAVLSGPNLFINNLNKTDNGTYRCEASNIVGKAHSDYMLYVYDPPTTIPPTTTTTTT  
TTTTTITITDATTTEPAVHDSRAGEEGTIGAVDHA VIGGVAVVVFAMLCLLILGRYFARHKGT  
FTHEAKGADDAADADTAIINAEGGQNNSEEKKEYFI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	50.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_997558</a>
Locus ID:	54725



[View online »](#)

UniProt ID:	<u>Q8R5M8</u>
RefSeq Size:	4387
Cytogenetics:	9 A5.3
RefSeq ORF:	1368
Synonyms:	2900073G06Rik; 3100001I08Rik; AI987920; BI2; Igsf4; Igsf4a; Necl2; RA175; RA175A; RA175B; RA175C
Summary:	<p>Mediates homophilic cell-cell adhesion in a Ca(2+)-independent manner. Also mediates heterophilic cell-cell adhesion with CADM3 and NECTIN3 in a Ca(2+)-independent manner. Acts as a tumor suppressor in non-small-cell lung cancer (NSCLC) cells. Interaction with CRTAM promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated rejection of tumors expressing CADM3 in vivo. May contribute to the less invasive phenotypes of lepodic growth tumor cells. In mast cells, may mediate attachment to and promote communication with nerves. CADM1, together with MITF, is essential for development and survival of mast cells in vivo. Acts as a synaptic cell adhesion molecule and plays a role in the formation of dendritic spines and in synapse assembly. May be involved in neuronal migration, axon growth, pathfinding, and fasciculation on the axons of differentiating neurons. May play diverse roles in the spermatogenesis including in the adhesion of spermatocytes and spermatids to Sertoli cells and for their normal differentiation into mature spermatozoa.[UniProtKB/Swiss-Prot Function]</p>