

## Product datasheet for TP522875

### Lamp2 (NM\_010685) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse lysosomal-associated membrane protein 2 (Lamp2), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR222875 representing NM_010685 <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	<p>MCLSPVKGAKLILIFLFLGAVQSNALIVNLTDSKGTCLYAEWEMNFTITYETTNQTNKTITIAVPDKATH DGSSCGDDRNSAKIMIQFGFAVSWAVNFTKEASHYSIHDIVLSYNTSDSTVFPGAVAKGVHTVKNPENFK VPLDVIFKCNVLTYNLTPVVQKYWGIHLQAFVQNGTVSKNEQVCEEDQPTTTPVAPIIHTTAPSTTTTLT PTSTPTPTPTPTVGNYSIRNGNTTCLLATMGLQLNITEEKVPFIFNINPATTNFTGSCQPQSAQLRLN NSQIKYLD FIFAVKNEKRFYLKEVNVYMYLANGSAFNISNKNLSFWDAPLGSSYMCNKEQVLSVSRAFQI NTFNLKVQPFNVTKGQYSTAQECSLDDDDTILPIIVGAGLSGLIIVIVIAYLIGRRKTYAGYQTL</p> <p><b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b></p>
Tag:	C-MYC/DDK
Predicted MW:	46.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_034815</a>
Locus ID:	16784
UniProt ID:	<a href="#">P17047</a>



[View online »](#)

RefSeq Size:	3603
Cytogenetics:	X 22.67 cM
RefSeq ORF:	1245
Synonyms:	CD107b; Lamp-2; Lamp-2a; Lamp-2b; Lamp-2c; Lamp II; LGP-B; Mac3
Summary:	<p>Plays an important role in chaperone-mediated autophagy, a process that mediates lysosomal degradation of proteins in response to various stresses and as part of the normal turnover of proteins with a long biological half-life (PubMed:10972293). Functions by binding target proteins, such as GAPDH and MLLT11, and targeting them for lysosomal degradation (By similarity). Required for the fusion of autophagosomes with lysosomes during autophagy (PubMed:27628032). Cells that lack LAMP2 express normal levels of VAMP8, but fail to accumulate STX17 on autophagosomes, which is the most likely explanation for the lack of fusion between autophagosomes and lysosomes (PubMed:27628032). Required for normal degradation of the contents of autophagosomes (PubMed:10972293, PubMed:12221139). Plays a role in lysosomal protein degradation in response to starvation (PubMed:27628032). Required for efficient MHCII-mediated presentation of exogenous antigens via its function in lysosomal protein degradation; antigenic peptides generated by proteases in the endosomal/lysosomal compartment are captured by nascent MHCII subunits. Is not required for efficient MHCII-mediated presentation of endogenous antigens (By similarity). [UniProtKB/Swiss-Prot Function]</p>