

## **Product datasheet for TP523008**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Tbk1 (NM 019786) Mouse Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse TANK-binding kinase 1 (Tbk1), with C-terminal

MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse Expression Host: HEK293T

**Expression cDNA Clone** >MR223008 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MQSTSNHLWLLSDILGQGATANVFRGRHKKTGDLYAVKVFNNISFLRPVDVQMREFEVLKKLNHKNIVKL FAIEEETTTRHKVLIMEFCPCGSLYTVLEEPSNAYGLPESEFLIVLRDVVGGMNHLRENGIVHRDIKPGN IMRVIGEDGQSVYKLTDFGAARELEDDEQFVSLYGTEEYLHPDMYERAVLRKDHQKKYGATVDLWSVGVT FYHAATGSLPFRPFEGPRRNKEVMYKIITGKPSGAISGVQKAENGPIDWSGDMPLSCSLSQGLQALLTPV LANILEADQEKCWGFDQFFAETSDVLHRMVIHVFSLQHMTAHKIYIHSYNTAAVFHELVYKQTKIVSSNQ ELIYEGRRLVLELGRLAQHFPKTTEENPIFVTSREQLNTVGLRYEKISLPKIHPRYDLDGDASMAKAVTG VVCYACRTASTLLLYQELMRKGVRWLVELVKDDYNETVHKKTEVVITLDFCIRNIEKTVKVYEKLMKVNL EAAELGEISDIHTKLLRLSSSQGTIESSLQDISSRLSPGGLLADTWAHQEGTHPRDRNVEKLQVLLNCIT EIYYQFKKDKAERRLAYNEEQIHKFDKQKLYYHATKAMSHFSEECVRKYEAFKDKSEEWMRKMLHLRKQL LSLTNQCFDIEEEVSKYQDYTNELQETLPQKMLAASGGVKHAMAPIYPSSNTLVEMTLGMKKLKEEMEGV

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

VKELAENNHILERFGSLTMDGGLRNVDCL

Tag: C-MYC/DDK
Predicted MW: 83.4 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C after receiving vials.





## Tbk1 (NM\_019786) Mouse Recombinant Protein - TP523008

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 062760

**Locus ID:** 56480

UniProt ID: Q9WUN2

RefSeq Size: 3031 Cytogenetics: 10 D2 RefSeq ORF: 2187

**Synonyms:** 1200008B05Rik; Al462036; AW048562

**Summary:** Serine/threonine kinase that plays an essential role in regulating inflammatory responses to

foreign agents (PubMed:10581243, PubMed:15210742, PubMed:15661922). Following activation of toll-like receptors by viral or bacterial components, associates with TRAF3 and TANK and phosphorylates interferon regulatory factors (IRFs) IRF3 and IRF7 as well as DDX3X (By similarity). This activity allows subsequent homodimerization and nuclear translocation of the IRFs leading to transcriptional activation of pro-inflammatory and antiviral genes including IFNA and IFNB (By similarity). In order to establish such an antiviral state, TBK1 form several different complexes whose composition depends on the type of cell and cellular stimuli (By similarity). Thus, several scaffolding molecules including FADD, TRADD, MAVS, AZI2, TANK or TBKBP1/SINTBAD can be recruited to the TBK1-containing-complexes (By similarity). Plays a key role in IRF3 activation: acts by first phosphorylating innate adapter proteins MAVS, TMEM173/STING and TICAM1 on their pLxIS motif, leading to recruitment of IRF3, thereby licensing IRF3 for phosphorylation by TBK1 (By similarity). Under particular conditions, functions as a NF-kappa-B effector by phosphorylating NF-kappa-B inhibitor alpha/NFKBIA, IKBKB or RELA to translocate NF-Kappa-B to the nucleus (By similarity). Restricts bacterial proliferation by phosphorylating the autophagy receptor OPTN/Optineurin on 'Ser-177', thus enhancing LC3 binding affinity and antibacterial autophagy (By similarity). Phosphorylates SMCR8 component of the C9orf72-SMCR8 complex, promoting autophagosome maturation (By similarity). Phosphorylates and activates AKT1 (By similarity). Seems to play a role in energy balance regulation by sustaining a state of chronic, low-grade inflammation in obesity, wich leads to a negative impact on insulin sensitivity (PubMed:23396211).[UniProtKB/Swiss-