

Product datasheet for TP524897

Cyp1a1 (NM_001136059) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse cytochrome P450, family 1, subfamily a, polypeptide 1 (Cyp1a1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR224897 protein sequence Red=Cloning site Green=Tags(s)

MPSMYGLPAFVSATELLAVTVFCLGFVWRATRTWVPKGLKTPPGPWGLPFIGHMLTVGKNPHLSLTRL
 SQYGDVLRIGSTPVVLSGLNTIKQALVRQGDDFKGRPDLYSFTLITNGKSMFNPDSGPVWAARRR
 LAQNALKSFSIASDPTSASSCYLEEHVSKEANYLVSKLQKVM AEVGHFDPYKYLVSVANVICAICFGQR
 YDHHDDQELLSIVNLSNEFGEVTGSGYPADFIPVLRYPNSSLDAFKDLNDKFYSFMKKLIKEHYRTFEKG
 HIRDITDSLIEHCQDRKLDENANVQLSDDKVITVLDLFGAGFDTVTTAISWSLMLVTNPRVQRKIQEE
 LDTVIGRDRQRLSDRPQLPYLEAFILETFRHSSFPFTIPHSTTRDTSLNFGFYIPKGCVCVFNQWQVNH
 DRELWGDPNEFRPERFLTPSGTLDKRLSEKVTFLGLGKRKCIGETIGRSEVFLFLAILLQQIEFKVSPGE
 KVDMTPTYGLTLKHARCEHFQVQMRSSGPQHLQA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	59.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001129531



[View online »](#)

Locus ID:	13076
UniProt ID:	P00184 , Q05A20
RefSeq Size:	2637
Cytogenetics:	9 31.34 cM
RefSeq ORF:	1575
Synonyms:	AHH; AHRR; CP11; Cyp1a2; CYPIA1; P450-1
Summary:	<p>A cytochrome P450 monooxygenase involved in the metabolism of various endogenous substrates, including fatty acids, steroid hormones and vitamins. Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase). Catalyzes the hydroxylation of carbon-hydrogen bonds. Exhibits high catalytic activity for the formation of hydroxysteroids from estrone (E1) and 17beta-estradiol (E2), namely 2-hydroxy E1 and E2, as well as D-ring hydroxylated E1 and E2 at the C15alpha and C16alpha positions. Displays different regioselectivities for polyunsaturated fatty acids (PUFA) hydroxylation. Catalyzes the epoxidation of double bonds of certain PUFA. Converts arachidonic acid toward epoxyeicosatrienoic acid (EET) regioisomers, 8,9-, 11,12-, and 14,15-EET, that function as lipid mediators in the vascular system. Displays an absolute stereoselectivity in the epoxidation of eicosapentaenoic acid (EPA) producing the 17(R),18(S) enantiomer. May play an important role in all-trans retinoic acid biosynthesis in extrahepatic tissues. Catalyzes two successive oxidative transformation of all-trans retinol to all-trans retinal and then to the active form all-trans retinoic acid. May also participate in eicosanoids metabolism by converting hydroperoxide species into oxo metabolites (lipoxygenase-like reaction, NADPH-independent).[UniProtKB/Swiss-Prot Function]</p>