

## Product datasheet for TP525184

### Prkcd (NM\_011103) Mouse Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse protein kinase C, delta (Prkcd), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR225184 representing NM_011103 <span style="color: red;">Red</span> =Cloning site <span style="color: green;">Green</span> =Tags(s)  MAPFLRISFNSYELGSLQVEDEASQPFCAVKMKEALSTERGKTLVQKKPTMYPEWKTTFDAHIYEGRVIQ IVLMRAAEDPVSEVTGVSVLAERCKKNGKAFFWLDLQPQAKVLMCVQYFLEDGDCKQSMRSEEEAKF P TMNRRGAIKQAKIHYIKNHEFIATFFGQPTFCVCKEFVWGLNKQGYKRCQCNAAIHKKCIDKIIGRCTG TATNSRDTIFQKERFNIDMPHRFKVYNYMSPTFCDHCGSLLWGLVKQGLKCEDCGMNVHHKCREKVAN LC GINQKLLAEALNQVTQRSSRKLDTTESVGIYQGFEKKPEVSGSDILDNNGTYGKIWEGSTRCTLENFTFQ KVLGKGSFGKVLLAELGKDKYFAIKCLKKDVLIDDDVECTMVEKRVLALAWESPFLTHLICTFQTKDH LFFVMEFLNGGDLMFHIQDKGRFELYRATFYAAEIIICGLQFLHSGKIIYRDLKLDNVMLDRDGHKIAF GMCKENIFGEGRASTFCGTPDYIAPEILQGLKYSFSVDWWSFGVLLYEMLIGQSPFHHGDDDELFEFESIRV DTPHYPRWITKESKDIMEKLFERDPDKRLGVTGNIRIHPFFKTINWSLLEKRVKVEPPFKPKVKSPSDYSN FDPEFLNEKPQLSFSKLNIDSMQEAHFGFSFVNPKFEQFLDI  <span style="color: red;">TR</span> <span style="color: green;">TRPLEQKLISEEDLAANDILDYKDDDDKV</span>
Tag:	C-MYC/DDK
Predicted MW:	78 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.


[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_035233</a>
Locus ID:	18753
UniProt ID:	<a href="#">P28867</a>
RefSeq Size:	2536
Cytogenetics:	14 18.82 cM
RefSeq ORF:	2022
Synonyms:	AI385711; D14Ert420e; Pkcd; PKCdelta; PKC[d]

**Summary:**

Calcium-independent, phospholipid- and diacylglycerol (DAG)-dependent serine/threonine-protein kinase that plays contrasting roles in cell death and cell survival by functioning as a pro-apoptotic protein during DNA damage-induced apoptosis, but acting as an anti-apoptotic protein during cytokine receptor-initiated cell death, is involved in tumor suppression, is required for oxygen radical production by NADPH oxidase and acts as positive or negative regulator in platelet functional responses. Negatively regulates B cell proliferation and also has an important function in self-antigen induced B cell tolerance induction. Upon DNA damage, activates the promoter of the death-promoting transcription factor BCLAF1/Btf to trigger BCLAF1-mediated p53/TP53 gene transcription and apoptosis. In response to oxidative stress, interact with and activate CHUK/IKKA in the nucleus, causing the phosphorylation of p53/TP53. In the case of ER stress or DNA damage-induced apoptosis, can form a complex with the tyrosine-protein kinase ABL1 which trigger apoptosis independently of p53/TP53. In cytosol can trigger apoptosis by activating MAPK11 or MAPK14, inhibiting AKT1 and decreasing the level of X-linked inhibitor of apoptosis protein (XIAP), whereas in nucleus induces apoptosis via the activation of MAPK8 or MAPK9. Upon ionizing radiation treatment, is required for the activation of the apoptosis regulators BAX and BAK, which trigger the mitochondrial cell death pathway. Can phosphorylate MCL1 and target it for degradation which is sufficient to trigger for BAX activation and apoptosis. Is required for the control of cell cycle progression both at G1/S and G2/M phases. Mediates phorbol 12-myristate 13-acetate (PMA)-induced inhibition of cell cycle progression at G1/S phase by up-regulating the CDK inhibitor CDKN1A/p21 and inhibiting the cyclin CCNA2 promoter activity. In response to UV irradiation can phosphorylate CDK1, which is important for the G2/M DNA damage checkpoint activation. Can protect glioma cells from the apoptosis induced by TNFSF10/TRAIL, probably by inducing increased phosphorylation and subsequent activation of AKT1. Can also act as tumor suppressor upon mitogenic stimulation with PMA or TPA. In N-formyl-methionyl-leucyl-phenylalanine (fMLP)-treated cells, is required for NCF1 (p47-phox) phosphorylation and activation of NADPH oxidase activity, and regulates TNF-elicited superoxide anion production in neutrophils, by direct phosphorylation and activation of NCF1 or indirectly through MAPK1/3 (ERK1/2) signaling pathways. May also play a role in the regulation of NADPH oxidase activity in eosinophil after stimulation with IL5, leukotriene B4 or PMA. In collagen-induced platelet aggregation, acts a negative regulator of filopodia formation and actin polymerization by interacting with and negatively regulating VASP phosphorylation. Downstream of PAR1, PAR4 and CD36/GP4 receptors, regulates differentially platelet dense granule secretion; acts as a positive regulator in PAR-mediated granule secretion, whereas it negatively regulates CD36/GP4-mediated granule release. Phosphorylates MUC1 in the C-terminal and regulates the interaction between MUC1 and beta-catenin. The catalytic subunit phosphorylates 14-3-3 proteins (YWHAB, YWHAZ and YWHAH) in a sphingosine-dependent fashion. Phosphorylates ELAVL1 in response to angiotensin-2 treatment (By similarity).[UniProtKB/Swiss-Prot Function]