

Product datasheet for **TP525488**

Fst (NM_008046) Mouse Recombinant Protein

Product data:

| | |
|---------------------------------------|--|
| Product Type: | Recombinant Proteins |
| Description: | Purified recombinant protein of Mouse follistatin (Fst), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug |
| Species: | Mouse |
| Expression Host: | HEK293T |
| Expression cDNA Clone or AA Sequence: | A DNA sequence from Mouse cDNA ORF Clone, MR225488, encoding Mouse full-length Fst. |
| Tag: | C-MYC/DDK |
| Predicted MW: | 38.2 kDa |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method |
| Purity: | > 80% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol |
| Note: | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. |
| Storage: | Store at -80°C after receiving vials. |
| Stability: | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. |
| RefSeq: | NP_032072 |
| Locus ID: | 14313 |
| UniProt ID: | P47931 |
| RefSeq Size: | 2337 |
| Cytogenetics: | 13 D2.2 |
| RefSeq ORF: | 1032 |
| Synonyms: | AL033346; D2Mgi5; FS |


[View online »](#)

Summary:

The protein encoded by this gene binds to and negatively regulates activin, as well as other members of the transforming growth factor beta family, and acts to prevent uncontrolled cellular proliferation. This protein also contains a heparin-binding sequence. It is expressed in many of the tissues in which activin is synthesized and is thought to clear activin from the circulation by attachment to the cell surface. Alternative splicing results in multiple transcript variants that encode multiple protein isoforms, including FST315 and FST288, that differ at their C-terminus. Another isoform, FST303 is thought to be produced by proteolytic cleavage of FST315. These isoforms differ in their localization and in their ability to bind heparin. While FST315 is a circulating protein, FST288 is tissue-bound, and FST303 is gonad-specific. While deletion of all isoforms results in embryonic lethality, expression of just FST288 is sufficient for embryonic development, but the resultant mice have fertility defects. [provided by RefSeq, Aug 2014]