

Product datasheet for TP525528

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Dmrt1 (NM_015826) Mouse Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Mouse doublesex and mab-3 related transcription factor 1

(Dmrt1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

Species: Mouse

Expression Host: HEK293T

Expression cDNA

>MR225528 representing NM_015826

Red=Cloning site Green=Tags(s)

Clone or AA Sequence:

MPNDDTFGKPSTPTEVPHAPGAPPQGKAGGYSKAAGAMAGAAGGSGAGGSGGASGSGPSGLGSGSKKSPR LPKCARCRNHGYASPLKGHKRFCMWRDCQCKKCSLIAERQRVMAAQVALRRQQAQEEELGISHPIPLPSA AELLVKRENNASNPCLMAENSSSAQPPPASTPTPAASEGRMVIQDIPAVTSRGHMENTSDLVSDPAYYSS FYQPSLFPYYNNLYNYPQYSMALSAESSSGEVGNSLGGSPVKNSLRSLPAPYVPAQTGNQWQMKTSESRH PVSSQYRMHSYYGPPSYLGQSMSQIFTFEEGPSYSEAKASVFSPPSSQDSGLVSLSSSSSPMSNESSKGVL

ECESASSEPSSYAVNQVLEEDEDE

SGPTRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-MYC/DDK

Predicted MW: 39.9 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C after receiving vials.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 056641

 Locus ID:
 50796

 UniProt ID:
 Q9QZ59





Dmrt1 (NM_015826) Mouse Recombinant Protein - TP525528

oogenesis.[UniProtKB/Swiss-Prot Function]

RefSeq Size: 2251

Cytogenetics: 19 B RefSeq ORF: 1122

Summary: Transcription factor that plays a key role in male sex determination and differentiation by

controlling testis development and male germ cell proliferation. Plays a central role in spermatogonia by inhibiting meiosis in undifferentiated spermatogonia and promoting mitosis, leading to spermatogonial development and allowing abundant and continuous production of sperm. Acts both as a transcription repressor and activator: prevents meiosis by restricting retinoic acid (RA)-dependent transcription and repressing STRA8 expression and promotes spermatogonial development by activating spermatogonial differentiation genes, such as SOHLH1. Also plays a key role in postnatal sex maintenance by maintaining testis determination and preventing feminization: represses transcription of female promoting genes such as FOXL2 and activates male-specific genes. May act as a tumor suppressor. May also play a minor role in