

Product datasheet for TP525846

Prkd1 (NM_008858) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse protein kinase D1 (Prkd1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR225846 representing NM_008858 Red =Cloning site Green =Tags(s)
	<p>MSVPPLLPPSPLLPAAAAVAAAAAALVPGSGPAPFPAPGAAPAGGISFHLQIGLSREPVLLLQDSSGDY SLAHVREMACSIVDQKFPECGFYGLYDKILLFRHDPASDNILQLVKIASDIQEGDLIEVLSASATFEDF QIRPHALFVHSYRAPAFCDHCGEMLWGLVRQGLKCEGCLNYHKRCAFKIPNNCSGVRRRRLSNVSLTGL GTVRTASAEFSTSVSPDEPLLSPVSPGFQKSPSEFIGREKRSNSQSYIGRPIQLDKLLMSKVKVPHTFV IHSYTRPTVCQFCKLLKGLFRQGLQCKDCRFNCHKRCAPKVPNNCLGEVTINGELLSPGAESDVMEEG SDDNDSERNISGLMDDMDEAMVQDTEMALAEGQSGGAEMQDPDADQEDSNRTISPSTSNNIPLMRVVQSVK HTKRRSSTVMKEGWMVHYTSKDTLRKRHYWRLDSKCITLQNDTGSRYEKEIPLSEILCLEPAKPSALTP VGATPHCFEITANVVVVYVGENVNPSSPPNNSVLPSPGIXPDVARMWEVAIQHALMPVIPKGSSVSGS SNSHKDISVSISVSNQCQIQENVDISTVYQIFPDEVLGSGQFGIVYGGKHKRKTGRDVAIKIIDKLRFPKQ ESQLRNEVAILQNLHHPGVNLECMFETPERVFMKLEHGDMLMILSSEKGRLEPHITKFLITQILVA LRHLHFKNIVHCDLKPENVLLASADPFQVKLCDFGFARIIGEKSFRRSVVGTAYLAPEVLRNKGYNRS LDMWSVGVIIYVSLSGTFPFNEDEDIHDQIQNAAFMYPNPWKEISHEAIDLINLLQVKMRKRYSDKT LSHPWLQDYQTWLDLRELECRIGERYITHESDSDSRWEQYAGEQGLQYPAHLISLSASHSDSPEAEEREMK ALSERSVIL</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	102.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol



[View online »](#)

Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_032884
Locus ID:	18760
UniProt ID:	Q62101
RefSeq Size:	3778
Cytogenetics:	12 B3
RefSeq ORF:	2757
Synonyms:	Pkcm; PKD; PKD1; Prkcm
Summary:	<p>Serine/threonine-protein kinase that converts transient diacylglycerol (DAG) signals into prolonged physiological effects downstream of PKC, and is involved in the regulation of MAPK8/JNK1 and Ras signaling, Golgi membrane integrity and trafficking, cell survival through NF-kappa-B activation, cell migration, cell differentiation by mediating HDAC7 nuclear export, cell proliferation via MAPK1/3 (ERK1/2) signaling, and plays a role in cardiac hypertrophy, VEGFA-induced angiogenesis, genotoxic-induced apoptosis and flagellin-stimulated inflammatory response. Phosphorylates the epidermal growth factor receptor (EGFR) on dual threonine residues, which leads to the suppression of epidermal growth factor (EGF)-induced MAPK8/JNK1 activation and subsequent JUN phosphorylation. Phosphorylates RIN1, inducing RIN1 binding to 14-3-3 proteins YWHAB, YWHAH and YWHAZ and increased competition with RAF1 for binding to GTP-bound form of Ras proteins (NRAS, HRAS and KRAS). Acts downstream of the heterotrimeric G-protein beta/gamma-subunit complex to maintain the structural integrity of the Golgi membranes, and is required for protein transport along the secretory pathway. In the trans-Golgi network (TGN), regulates the fission of transport vesicles that are on their way to the plasma membrane. May act by activating the lipid kinase phosphatidylinositol 4-kinase beta (PI4KB) at the TGN for the local synthesis of phosphorylated inositol lipids, which induces a sequential production of DAG, phosphatidic acid (PA) and lyso-PA (LPA) that are necessary for membrane fission and generation of specific transport carriers to the cell surface. Under oxidative stress, is phosphorylated at Tyr-469 via SRC-ABL1 and contributes to cell survival by activating IKK complex and subsequent nuclear translocation and activation of NFKB1. Involved in cell migration by regulating integrin alpha-5/beta-3 recycling and promoting its recruitment in newly forming focal adhesion. In osteoblast differentiation, mediates the bone morphogenetic protein 2 (BMP2)-induced nuclear export of HDAC7, which results in the inhibition of HDAC7 transcriptional repression of RUNX2. In neurons, plays an important role in neuronal polarity by regulating the biogenesis of TGN-derived dendritic vesicles, and is involved in the maintenance of dendritic arborization and Golgi structure in hippocampal cells. May potentiate mitogenesis induced by the neuropeptide bombesin or vasopressin by mediating an increase in the duration of MAPK1/3 (ERK1/2) signaling, which leads to accumulation of immediate-early gene products including FOS that stimulate cell cycle</p>

progression. Plays an important role in the proliferative response induced by low calcium in keratinocytes, through sustained activation of MAPK1/3 (ERK1/2) pathway. Downstream of novel PKC signaling, plays a role in cardiac hypertrophy by phosphorylating HDAC5, which in turn triggers XPO1/CRM1-dependent nuclear export of HDAC5, MEF2A transcriptional activation and induction of downstream target genes that promote myocyte hypertrophy and pathological cardiac remodeling. Mediates cardiac troponin I (TNNI3) phosphorylation at the PKA sites, which results in reduced myofilament calcium sensitivity, and accelerated crossbridge cycling kinetics. The PRKD1-HDAC5 pathway is also involved in angiogenesis by mediating VEGFA-induced specific subset of gene expression, cell migration, and tube formation. In response to VEGFA, is necessary and required for HDAC7 phosphorylation which induces HDAC7 nuclear export and endothelial cell proliferation and migration. During apoptosis induced by cytarabine and other genotoxic agents, PRKD1 is cleaved by caspase-3 at Asp-378, resulting in activation of its kinase function and increased sensitivity of cells to the cytotoxic effects of genotoxic agents. In epithelial cells, is required for transducing flagellin-stimulated inflammatory responses by binding and phosphorylating TLR5, which contributes to MAPK14/p38 activation and production of inflammatory cytokines.