

Product datasheet for TP525859

Gnas (NM_201616) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse GNAS (guanine nucleotide binding protein, alpha stimulating) complex locus (Gnas), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR225859 representing NM_201616 Red =Cloning site Green =Tags(s) MGCLGNSKTEDQRNEEKAQREANKKIEKQLQKDKQVYRATHRLLLLGAGESGKSTIVKQMRILHVNGFN G EGGEEDPQAARSNSDGEKATKVQDIKNNLKEAIETIVAAMSNLVPVELANPENQFRVDYILSVMNVPNF DFPPEFYEHAKALWEDEGVRACYERSNEYQLIDCAQYFLDKIDVIKQADYVPSDQDLLRCRVLTSGIFET KFQVDKVNFMFDVGGQRDERRKWIQCFNDVTAIIFVASSSYNMVIREDNQTNRLQEALNLFKSIWNN R WLRTISVILFLNKQDLLAEKVLGKSKIETYFPEFARYTTPEDATPEPGEDPRVTRAKYFIRDEFRLRIST ASGDGRHYCYPHFTCAVDTENIRRVFNDCRDIIQRMHLRQYELL SGPTRTRPLE QKLISEEDLA NDILDYKDDDDKV
Tag:	C-MYC/DDK
Predicted MW:	46.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.


[View online »](#)

RefSeq:	<u>NP_963910</u>
Locus ID:	14683
UniProt ID:	<u>P63094</u>
RefSeq Size:	1762
Cytogenetics:	2 97.89 cM
RefSeq ORF:	1182
Synonyms:	5530400H20Rik; A930027G11Rik; C130027O20Rik; G; Ga; Galphas; Gn; Gnas1; Gnasxl; GPSA; Gs-; Gs-alpha; Gsa; GSP; N; Nes; Nesp; Nesp55; Nespl; Oed; Oed-Sml; Oedsml; P; P1; P2; P3; PHP1A; PHP1B; POH; SCG; SCG6; XL
Summary:	<p>This locus has a highly complex imprinted expression pattern. It gives rise to maternally, paternally, and biallelically expressed transcripts that are derived from four alternative promoters and 5' exons. Some transcripts contain a differentially methylated region (DMR) at their 5' exons, which is commonly found in imprinted genes and correlates with transcript expression. This gene has an antisense transcript. One of the transcripts produced from this locus, and the antisense transcript, are both paternally expressed noncoding RNAs, and may regulate imprinting in this region. In addition, one of the transcripts contains a second overlapping ORF, which encodes a structurally unrelated protein - Alex. Alternative splicing of downstream exons is also observed, which results in different forms of the stimulatory G-protein alpha subunit, a key element of the classical signal transduction pathway linking receptor-ligand interactions with the activation of adenylyl cyclase and a variety of cellular responses. Additional transcript variants have been found for this gene, but the full-length nature and/or biological validity of some variants have not been determined. [provided by RefSeq, Jun 2015]</p>