

Product datasheet for TP526558

Cbln1 (NM_019626) Mouse Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse cerebellin 1 precursor protein (Cbln1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR226558 representing NM_019626 Red=Cloning site Green=Tags(s)

MLGWELLLLGTAWLAGPARGQNETEPIVLEGKCLWCDSNPTSDPTGTALGISVRSKSAKVAFSAIRST
NHEPSEMSNRMTIIYFDQVLVNIIGNNFDSESTFIAPRKGISFNHFWKVVYNRQTIQVSLMLNGWPVIS
AFAGDQDVTREAASNGVLIQMEKGDRAVYKLERGNLMGGWKYSTFSGLVFPL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-MYC/DDK
Predicted MW:	21.6 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C after receiving vials.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_062600</u>
Locus ID:	12404
UniProt ID:	<u>Q9R171</u> , <u>Q7TNF5</u>
RefSeq Size:	2345
Cytogenetics:	8 42.16 cM



[View online »](#)

RefSeq ORF: 579

Synonyms: AI323299

Summary: Required for synapse integrity and synaptic plasticity. During cerebellar synapse formation, essential for the matching and maintenance of pre- and post-synaptic elements at parallel fiber-Purkinje cell synapses, the establishment of the proper pattern of climbing fiber-Purkinje cell innervation, and induction of long-term depression at parallel fiber-Purkinje cell synapses (PubMed:16234806). Plays a role as a synaptic organizer that acts bidirectionally on both pre- and post-synaptic components (PubMed:20395510). On the one hand induces accumulation of synaptic vesicles in the pre-synaptic part by binding with NRXN1 and in other hand induces clustering of GRID2 and its associated proteins at the post-synaptic site through association of GRID2 (PubMed:21410790). NRXN1-CBLN1-GRID2 complex directly induces parallel fiber protrusions that encapsulate spines of Purkinje cells leading to accumulation of GRID2 and synaptic vesicles (PubMed:23141067). Required for CBLN3 export from the endoplasmic reticulum and secretion (PubMed:17030622, PubMed:17331201). NRXN1-CBLN1-GRID2 complex mediates the D-Serine-dependent long term depression signals and AMPA receptor endocytosis (By similarity).[UniProtKB/Swiss-Prot Function]