

## Product datasheet for TP526792

### Nfkb1 (NM\_008689) Mouse Recombinant Protein

#### Product data:


Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse nuclear factor of kappa light polypeptide gene enhancer in B cells 1, p105 (Nfkb1), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR226792 protein sequence <span style="color: red;">Red</span> =Cloning site <span style="color: green;">Green</span> =Tags(s)  MADDDPYGTGQM FHLNTALTHSIFNAELYSPEIPLSTDGPYLQILEQPKQGRGFRFRVYCEGSPSHGGLPGA SSEKNKKSYPQVKICNYVGPAKVIVQLVTNGKNIHLHAHSLVGKHCEDGVCTVTAGPKDMVVG FANLGIL HVTKKKVFETLEARMTEACIRGYNPGLLVHSDLAYLQAEGGGDRQLTDREKEIRQAAVQQTKEMDLSVV RLMFTAFLPDSTGSFTRRLEPVVSDAIYDSKAPNASNLKIVRMDRTAGCVTGGEIYLLCDKVQKDDIQI RFYEEEEENGWEGFGDFSPD VHRQFAIVFKTPKYKDVNITKPASVFVQLRRKSDLETSEPKPFLYPE IKDKEEVQRKRQKLMPNFSDSFGGGSGAGAGGGGMFGSGGGGGSTGSPGPYGYSNYGFPPYGGITFHP G VTKSNA GVTHTGINTKFKNGPKDCAKSDDEESLTLPKETEGEGPSLPMACTKTEPIALASTMEDKEQDM GFQDNL FLEKALQLARRHANALFDYAVTG DVKMLLAVQRHLTAVQDENGDSVLHLAIHLHAQLVRDLLE VTSG LISDDIINMRNDLYQTPLHLAVITKQEDVVEDLLRVGADLSLLDRWGN SVLHLAAKEGHDRILSIL LKSRKAAPLIDHPNGEGLNAIHIAVMSNSLPCLLLLVAAGAEVNAQE QKSGRTALHLAVEYDNISLAGCL LLEGDAHV DSTTYDGTTP L HIAAGRGSTRLAALLKAAGADPLVENFEPLYDLDDSWEKAGEDEGVVPGTT PLDMAANWQVFDILNGKPYEPVFTSDDILPQGDMKQLTEDTRLQLCKLLEIPDPDKNWATLAQKLGLGIL NNAFRLSPAPSKT LMDNYEVSGGTIKELMEALQ QMGYTEAIEVIQA AFRTPATTASSPVTTA QVHCLPLS SSSTRQHIDELRDS DSVCDSGVETSFRKLSFTESLTGDSPLLSLNKMPHGYGQEGPIEGKI  <span style="color: red;">TR</span> <span style="color: green;">TRPLEQKLISEEDLAANDILDYKDDDDKV</span>
Tag:	C-MYC/DDK
Predicted MW:	105.6 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol


[View online »](#)

<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C after receiving vials.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<u><a href="#">NP_032715</a></u>
<b>Locus ID:</b>	18033
<b>UniProt ID:</b>	<u><a href="#">P25799</a></u>
<b>RefSeq Size:</b>	4128
<b>Cytogenetics:</b>	3 62.82 cM
<b>RefSeq ORF:</b>	2913
<b>Synonyms:</b>	NF-kappaB; NF-kappaB1; NF-KB1; p50; p50/p105; p105

**Summary:**

NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and RelB-p50 complexes are transcriptional activators. The NF-kappa-B p50-p50 homodimer is a transcriptional repressor, but can act as a transcriptional activator when associated with BCL3. NFKB1 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p105 and generation of p50 by a cotranslational processing. The proteasome-mediated process ensures the production of both p50 and p105 and preserves their independent function, although processing of NFKB1/p105 also appears to occur post-translationally. p50 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. Plays a role in the regulation of apoptosis. Isoform 5, isoform 6 and isoform 7 act as inhibitors of transactivation of p50 NF-kappa-B subunit, probably by sequestering it in the cytoplasm. Isoform 3 (p98) (but not p84 or p105) acts as a transactivator of NF-kappa-B-regulated gene expression. In a complex with MAP3K8, NFKB1/p105 represses MAP3K8-induced MAPK signaling; active MAP3K8 is released by proteasome-dependent degradation of NFKB1/p105.[UniProtKB/Swiss-Prot Function]

**Product images:**

Purified recombinant protein Nfkb1 was analyzed by SDS-PAGE gel and Coomossie Blue Staining.