

## Product datasheet for TP527248

### Src (NM\_009271) Mouse Recombinant Protein

#### Product data:

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Mouse Rous sarcoma oncogene (Src), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug

**Species:** Mouse

**Expression Host:** HEK293T

**Expression cDNA Clone or AA Sequence:** >MR227248 representing NM\_009271  
**Red**=Cloning site **Green**=Tags(s)

MGSNKS PKDASQRRRSLEPSENVHGAGGAFPASQTPSKPASADGHRGPSAAFVPPAAEPKLFGGFNSSD  
TVTSPQRAGPLAGGVTTFFVALYDYESRTETDLSFKKGERLQIVNNTKVDVREGDWWLAHSLSTGQTGYI  
PSNYVAPSDSIQAEWYFGKITRRESERLLLNAENPRGTFVRESETTKGAYCLSVSDFDNAKGLNVKHY  
KIRKLDGGFYITSRTQFNSLQQLVAYYSKHADGLCHRLTTVCPSTSKPQTQGLAKDAWEIPRESLRLEVK  
LGQGCFGEVWMGTWNGTTRVAIKTLKPGTMSPEAFLQEAQVMKKLRHEKLVQLYAVVSEPIYIVTEYMN  
KGSLLDFLKGETGKYLRPLQLVDMASQIASGMAYVERMNYVHRDLRAANILVGENLVCKVADFGLARLIE  
DNEYTARQGAKFPIKWTAPEAALYGRFTIKSDVWSFGILLTELTTKGRVPYPMNVNREVLDQVERGYRMP  
CPPECPESLHDLMCQCWRKEPEERPTFEYLQAFLEDYFTSTEPQYQPGENL

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

**Tag:** C-MYC/DDK

**Predicted MW:** 61.1 kDa

**Concentration:** >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

**Storage:** Store at -80°C after receiving vials.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** [NP\\_033297](#)



[View online »](#)

Locus ID:	20779
UniProt ID:	<a href="#">P05480</a> , <a href="#">F8WI90</a>
RefSeq Size:	3887
Cytogenetics:	2 78.35 cM
RefSeq ORF:	1623
Synonyms:	AW259666; pp60c-src
Summary:	<p>Non-receptor protein tyrosine kinase which is activated following engagement of many different classes of cellular receptors including immune response receptors, integrins and other adhesion receptors, receptor protein tyrosine kinases, G protein-coupled receptors as well as cytokine receptors. Participates in signaling pathways that control a diverse spectrum of biological activities including gene transcription, immune response, cell adhesion, cell cycle progression, apoptosis, migration, and transformation. Due to functional redundancy between members of the SRC kinase family, identification of the specific role of each SRC kinase is very difficult. SRC appears to be one of the primary kinases activated following engagement of receptors and plays a role in the activation of other protein tyrosine kinase (PTK) families. Receptor clustering or dimerization leads to recruitment of SRC to the receptor complexes where it phosphorylates the tyrosine residues within the receptor cytoplasmic domains. Plays an important role in the regulation of cytoskeletal organization through phosphorylation of specific substrates such as AFAP1. Phosphorylation of AFAP1 allows the SRC SH2 domain to bind AFAP1 and to localize to actin filaments. Cytoskeletal reorganization is also controlled through the phosphorylation of cortactin (CTTN) (Probable). When cells adhere via focal adhesions to the extracellular matrix, signals are transmitted by integrins into the cell resulting in tyrosine phosphorylation of a number of focal adhesion proteins, including PTK2/FAK1 and paxillin (PXN) (By similarity). In addition to phosphorylating focal adhesion proteins, SRC is also active at the sites of cell-cell contact adherens junctions and phosphorylates substrates such as beta-catenin (CTNNB1), delta-catenin (CTNND1), and plakoglobin (JUP). Another type of cell-cell junction, the gap junction, is also a target for SRC, which phosphorylates connexin-43 (GJA1). SRC is implicated in regulation of pre-mRNA-processing and phosphorylates RNA-binding proteins such as KHDRBS1 (Probable). Also plays a role in PDGF-mediated tyrosine phosphorylation of both STAT1 and STAT3, leading to increased DNA binding activity of these transcription factors (PubMed:9344858). Involved in the RAS pathway through phosphorylation of RASA1 and RASGRF1. Plays a role in EGF-mediated calcium-activated chloride channel activation (By similarity). Required for epidermal growth factor receptor (EGFR) internalization through phosphorylation of clathrin heavy chain (CLTC and CLTCL1) at 'Tyr-1477'. Involved in beta-arrestin (ARRB1 and ARRB2) desensitization through phosphorylation and activation of GRK2, leading to beta-arrestin phosphorylation and internalization. Has a critical role in the stimulation of the CDK20/MAPK3 mitogen-activated protein kinase cascade by epidermal growth factor (Probable). Might be involved not only in mediating the transduction of mitogenic signals at the level of the plasma membrane but also in controlling progression through the cell cycle via interaction with regulatory proteins in the nucleus (By similarity). Plays an important role in osteoclastic bone resorption in conjunction</p>

with PTK2B/PYK2. Both the formation of a SRC-PTK2B/PYK2 complex and SRC kinase activity are necessary for this function. Recruited to activated integrins by PTK2B/PYK2, thereby phosphorylating CBL, which in turn induces the activation and recruitment of phosphatidylinositol 3-kinase to the cell membrane in a signaling pathway that is critical for osteoclast function (PubMed:14739300). Promotes energy production in osteoclasts by activating mitochondrial cytochrome C oxidase (PubMed:12615910). Phosphorylates DDR2 on tyrosine residues, thereby promoting its subsequent autophosphorylation. Phosphorylates RUNX3 and COX2 on tyrosine residues, TNK2 on 'Tyr-284' and CBL on 'Tyr-738'. Enhances DDX58/RIG-I-elicited antiviral signaling. Phosphorylates PDPK1 at 'Tyr-9', 'Tyr-373' and 'Tyr-376'. Phosphorylates BCAR1 at 'Tyr-226'. Phosphorylates CBLC