

## Product datasheet for TP527409

### Igf1r (NM\_010513) Mouse Recombinant Protein


#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Mouse insulin-like growth factor I receptor (Igf1r), with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug
Species:	Mouse
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>MR227409 representing NM_010513 Red=Cloning site Green=Tags(s)
	<p>MKSGSGGGSPTSLWGLVFLSAALSLWPTSGEICPGIDIRNDYQQLKRENCTVIEGFLHILLISKAEDY RSYRFPKLTVITEYLLFRVAGLESLGDLFPNLTVIRGWKLFYNYALVIFEMTNLKDIGLYNLRNITRGA IRIEKNADLCYLSTIDWSLILDAVSNYIVGNKPPKECGDLCPTLEEKPMCEKTTINNEYNYRCWTTNR CQKMCPSVCGKRAC TENNECCHPECLGSCHTPDDNTT CVACRHYYKGVCPACPPGTYRFEGWRCVDRD FCANIPNAESSDSDFVIHDDCEM QECPSGFIRNSTQSMYCIPCEGPCPKVCGDEEKTKTIDSVTSAQM LQGCTILKGNLLINIRRGNNIASELENFMGLIEVVTGYVKIRHSHALVLSFLKNLRLILGEEQLEGNYS FYVLDNQNQLQWLWDWNHRNLT VRSKGMYFAFNPKLCVSEIYRMEEVTGTKGRQSKGDINTRNNGERASCE SDVLRFTSTTTWKNRIITWHRYRPPDYRDLISFTVYYKEAPFKNVTEYDGDQACGSNSWNMVDVLDLPPN KEGEPGILLHGLKPWTQYAVYKAVTLTMVENDHIRGAKSEILYIRTNASVPSIPLDVL SASNSSSQLIV KWNPP TLPNGNLSYIVRWQRQPQDGYLRHNYCSKDKIPIRKYADGTIDVEEVTENPKTEVCGDGKGPC CACPKTEAEKQAEKEEA EYRKVFENFLHNSIFVPRPERRRRDVMQVANTTMSRSRNTTVADTYNITDPE EFETEYPPFESRVDNKERTVISNLRPFTLYRIDIHSCNHEAEKLGCSASN FVFARTMPAEGADDIPGPVT WEPRPENSIFLKWPEPENPNGLILMYEIKYGSQVEDQRECVSRQEYRKYGGAKLNRLNPGNYTARIQATS LSGNGSWTDPVFFYVPAKTTYENFMHLI ALPVAILLVGGGLVIMLYVFHRKRNSRLGNGVLYASV NPE YFSAADVVPDEWEVAREKITM NRELQGGSGFMVYEGVAKGVKDEPETRVAIKTVNEAASMRERIEFLN EASVMKEFNCHHWVRL LGVVSQGQPTLVIMELMTRGDLKSYLRSLRPEVEQNNLVLIPPSLSKMIQMAGE IADGMAYLNANKFVHRDLAARNCMVAEDFTVKIGDFGMTRDIYETDYRKGKGLLPVRWMSPELKDGV FTTHSDVWSFGVVLWEIATLAEQPYQGLSNEQVLR FVMEGGLLDKPDNCPDMLFELMRMCWQYNPKMRPS FLEIIGSIKDEMEPSFQEV SFYSEENKPEPEELEMENMESVPLDPSASSASLPLPERHSGHKAENG PGPGVLVLRASFDERQPYAHMNGGRANER ALPLQSSTC</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Tag:	C-MYC/DDK
Predicted MW:	155.7 kDa



[View online »](#)

<b>Concentration:</b>	>0.05 µg/µL as determined by microplate BCA method
<b>Purity:</b>	> 80% as determined by SDS-PAGE and Coomassie blue staining
<b>Buffer:</b>	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C after receiving vials.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_034643</a>
<b>Locus ID:</b>	16001
<b>UniProt ID:</b>	<a href="#">Q60751</a> , <a href="#">E9QNX9</a>
<b>RefSeq Size:</b>	11978
<b>Cytogenetics:</b>	7 37.27 cM
<b>RefSeq ORF:</b>	4107
<b>Synonyms:</b>	A330103N21Rik; CD221; D930020L01; hyft; IGF-1R
<b>Summary:</b>	<p>Receptor tyrosine kinase which mediates actions of insulin-like growth factor 1 (IGF1). Binds IGF1 with high affinity and IGF2 and insulin (INS) with a lower affinity. The activated IGF1R is involved in cell growth and survival control. IGF1R is crucial for tumor transformation and survival of malignant cell. Ligand binding activates the receptor kinase, leading to receptor autophosphorylation, and tyrosines phosphorylation of multiple substrates, that function as signaling adapter proteins including, the insulin-receptor substrates (IRS1/2), Shc and 14-3-3 proteins. Phosphorylation of IRSs proteins lead to the activation of two main signaling pathways: the PI3K-AKT/PKB pathway and the Ras-MAPK pathway. The result of activating the MAPK pathway is increased cellular proliferation, whereas activating the PI3K pathway inhibits apoptosis and stimulates protein synthesis. Phosphorylated IRS1 can activate the 85 kDa regulatory subunit of PI3K (PIK3R1), leading to activation of several downstream substrates, including protein AKT/PKB. AKT phosphorylation, in turn, enhances protein synthesis through mTOR activation and triggers the antiapoptotic effects of IGF1R through phosphorylation and inactivation of BAD. In parallel to PI3K-driven signaling, recruitment of Grb2/SOS by phosphorylated IRS1 or Shc leads to recruitment of Ras and activation of the ras-MAPK pathway. In addition to these two main signaling pathways IGF1R signals also through the Janus kinase/signal transducer and activator of transcription pathway (JAK/STAT). Phosphorylation of JAK proteins can lead to phosphorylation/activation of signal transducers and activators of transcription (STAT) proteins. In particular activation of STAT3, may be essential for the transforming activity of IGF1R. The JAK/STAT pathway activates gene transcription and may be responsible for the transforming activity. JNK kinases can also be activated by the IGF1R. IGF1 exerts inhibiting activities on JNK activation via phosphorylation and inhibition of MAP3K5/ASK1, which is able to directly associate with the IGF1R (By similarity). When present in a hybrid receptor with INSR, binds IGF1 (By similarity).[UniProtKB/Swiss-Prot Function]</p>

**Product images:**

Purified recombinant protein Igf1r was analyzed by SDS-PAGE gel and Coomassie Blue Staining.