

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TP700144

## Eph receptor A7 (EPHA7) (NM\_004440) Human Recombinant Protein

## **Product data:**


Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of human EPH receptor A7 (EPHA7), with C-terminal DDK/His tag, expressed in human cells, 20 μg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	A DNA sequence from TrueORF clone, RC600044, encoding the region (Gln28-Val555) of human EPHA7
Tag:	C-DDK/His
Predicted MW:	62 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	PBS, pH 7.4, 10% glycerol
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP 004431</u>
Locus ID:	2045
	2045
UniProt ID:	<u>Q15375</u>
UniProt ID: RefSeq Size:	
	<u>Q15375</u>
RefSeq Size:	<u>Q15375</u> 6644
RefSeq Size: Cytogenetics:	<u>Q15375</u> 6644 6q16.1



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Eph receptor A7 (EPHA7) (NM_004440) Human Recombinant Protein – TP700144
Summary:	This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Increased expression of this gene is associated with multiple forms of carcinoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2013]
Protein Families:	Druggable Genome, Protein Kinase, Transmembrane
Protein Pathway	s: Axon guidance

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US