

## **Product datasheet for TP700158**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## ROR1 (NM\_005012) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Homo sapiens receptor tyrosine kinase-like orphan receptor

1 (ROR1), transcript variant 1, residue 30-406 aa, expressed in HEK293 cells.

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

Predicted MW:

A DNA sequence from TrueORF clone, RC214967, encoding the extracellular domain (Gln30 -

Tyr406) of human ROR1, transcript variant 1

Tag: C-DDK/His

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** PBS, pH 7.4, 10% glycerol

45.1 kDa

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 005003

 Locus ID:
 4919

 UniProt ID:
 Q01973

 RefSeq Size:
 3358

Cytogenetics: 1p31.3 RefSeq ORF: 2811

**Synonyms:** dJ537F10.1; NTRKR1





**Summary:** 

This gene encodes a receptor tyrosine kinase-like orphan receptor that modulates neurite growth in the central nervous system. The encoded protein is a glycosylated type I membrane protein that belongs to the ROR subfamily of cell surface receptors. It is a pseudokinase that lacks catalytic activity and may interact with the non-canonical Wnt signalling pathway. This gene is highly expressed during early embryonic development but expressed at very low levels in adult tissues. Increased expression of this gene is associated with B-cell chronic lymphocytic leukaemia. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jun 2012]

**Protein Families:** 

Druggable Genome, Protein Kinase, Transmembrane

## **Product images:**

